

Anti-PNPT1 antibody ab157109

[3 References](#) [2 图像](#)

概述

产品名称	Anti-PNPT1抗体
描述	兔多克隆抗体to PNPT1
宿主	Rabbit
经测试应用	适用于: WB, IP
种属反应性	与反应: Human 预测可用于: Mouse, Rat, Rabbit, Horse, Dog, Pig, Chimpanzee, Ferret, Macaque monkey, Rhesus monkey, Common marmoset 
免疫原	Synthetic peptide corresponding to Human PNPT1 aa 700-800. Database link: Q8TCS8  Run BLAST with  Run BLAST with
阳性对照	293T, HeLa and Jurkat whole cell lysate (ab7899).
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	pH: 7 Preservative: 0.09% Sodium azide Constituent: 99% Tris citrate/phosphate pH 7 to 8
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用 ab157109 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
WB		1/1000 - 1/5000. Predicted molecular weight: 86 kDa.
IP		Use at 2-10 µg/mg of lysate.

靶标

相关性 PNPT1 is a subunit of the exosome complex, which is involved in 3-prime-to-5-prime exoribonuclease activity for RNA processing and degradation. It contains one KH domain and one S1 motif domain.

细胞定位 Mitochondrion

图片



All lanes : Anti-PNPT1 antibody (ab157109) at 0.4 µg/ml

Lane 1 : 293T cell lysate at 50 µg

Lane 2 : 293T cell lysate at 15 µg


Lane 3 : HeLa cell lysate at 50 µg

Lane 4 : Jurkat cell lysate at 50 µg

Developed using the ECL technique.

Predicted band size: 86 kDa

Exposure time: 3 minutes



Detection of PNPT1 by Western blot of immunoprecipitate.
 Immunoprecipitation of HeLa whole cell lysate at 6 µg/mg lysate (1 mg/IP; 20% of IP loaded/lane), using the following:
 Lane 1 : IP with an alternative Rabbit Anti-PNPT1 antibody.
 Lane 2 : IP with ab157109.
 Lane 3 : Control IgG.
 The subsequent Western blot of immunoprecipitates was done using ab157109 at 1 µg/ml.
 Detection: Chemiluminescence with exposure time of 30 seconds.

All lanes : Anti-PNPT1 antibody (ab157109)

All lanes : 293T whole cell lysate

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors