

Anti-Podoplanin antibody [RTD4E10] - BSA and Azide free ab11936

★★★★★ [12 Abreviews](#) [81 References](#) [2 图像](#)

概述

产品名称	Anti-Podoplanin抗体[RTD4E10] - BSA and Azide free
描述	叙利亚仓鼠单克隆抗体[RTD4E10] to Podoplanin - BSA and Azide free
宿主	Syrian hamster
特异性	The monoclonal antibody will detect podoplanin on the surface of mouse lymphatic endothelial cells and some epithelial cell types by immunostaining or immunohistochemistry. This antibody is designed to detect podoplanin, which is also referred to as gp38 (mouse and rat) and gp36 (human).
经测试应用	适用于: IHC-P
种属反应性	与反应: Mouse 不与反应: Human
免疫原	Tissue, cells or virus. This information is proprietary to Abcam and/or its suppliers.
阳性对照	Mouse normal skin FFPE tissue sections (IHC-P)
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Constituent: PBS
无载体	是
纯度	Protein G purified
纯化说明	Purified from cell culture supernatant.

克隆	单克隆
克隆编号	RTD4E10
同种型	IgG1

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab11936于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (6)	1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

靶标

功能	May be involved in cell migration and/or actin cytoskeleton organization. When expressed in keratinocytes, induces changes in cell morphology with transfected cells showing an elongated shape, numerous membrane protrusions, major reorganization of the actin cytoskeleton, increased motility and decreased cell adhesion. Required for normal lung cell proliferation and alveolus formation at birth. Induces platelet aggregation. Does not have any effect on folic acid or amino acid transport. Does not function as a water channel or as a regulator of aquaporin-type water channels.
组织特异性	Highly expressed in placenta, lung, skeletal muscle and brain. Weakly expressed in brain, kidney and liver. In placenta, expressed on the apical plasma membrane of endothelium. In lung, expressed in alveolar epithelium. Up-regulated in colorectal tumors and expressed in 25% of early oral squamous cell carcinomas.
序列相似性	Belongs to the podoplanin family.
翻译后修饰	Extensively O-glycosylated. Contains sialic acid residues. O-glycosylation is necessary for platelet aggregation activity. The N-terminus is blocked.
细胞定位	Membrane. Cell projection > filopodium membrane. Cell projection > lamellipodium membrane. Cell projection > microvillus membrane. Cell projection > ruffle membrane. Localized to actin-rich microvilli and plasma membrane projections such as filopodia, lamellipodia and ruffles.

图片




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Syrian hamster monoclonal [RTD4E10] to Podoplanin - BSA and Azide free (ab11936)

IHC image of Podoplanin staining in a section of formalin-fixed paraffin-embedded normal mouse skin. The section was pre-treated using pressure cooker heat mediated antigen retrieval with sodium citrate buffer (pH6). Non-specific protein-protein interactions were then blocked in TBS containing 0.025% (v/v) Triton X-100, 0.3M (w/v) glycine and 1% (w/v) BSA for 1h at room temperature. The section was then incubated overnight at +4°C in TBS containing 0.025% (v/v) Triton X-100 and 1% (w/v) BSA with ab11936 at 1/250. The secondary antibody (shown in green) was **ab180063** at 1/500 (Goat Anti-Syrian hamster IgG H&L (Alexa Fluor® 488) and counterstained using **ab190573**, Rabbit monoclonal [EP1332Y] to alpha Tubulin - Microtubule Marker (Alexa Fluor® 647), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue). The section was then mounted using Fluoromount®.

The negative control (inset) is a secondary-only assay to demonstrate low non-specific binding of the secondary antibody.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

For other IHC staining systems (automated and non-automated), customers should optimize variable parameters such as antigen retrieval conditions, antibody concentrations and incubation times.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Syrian hamster monoclonal [RTD4E10] to Podoplanin - BSA and Azide free (ab11936)

Image courtesy of an anonymous Abreview.

ab11936 staining Podoplanin in murine skin tissue sections by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). Tissue was fixed in paraformaldehyde and an enzymatic antigen retrieval step was performed using Protein K. Samples were then blocked with 5% serum for 30 minutes at room temperature and then incubated with ab11936 at a 1/400 dilution for 16 hours at 4°C. The secondary used was a biotin conjugated goat anti-hamster IgG used at a 1/500 dilution.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors