


Anti-RhoGDI antibody [EPR3773] ab133248

敲除验证 重组 RabMAb

★★★★★ 1 Abreviews 7 References 6 图像

概述

产品名称	Anti-RhoGDI抗体[EPR3773]
描述	兔单克隆抗体[EPR3773] to RhoGDI
宿主	Rabbit
经测试应用	适用于: WB, IHC-P 不适用于: Flow Cyt or IP
种属反应性	与反应: Mouse, Human 预测可用于: Rat 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HEK293T, Jurkat, HeLa and NIH3T3 cell lysates. IHC-P: Human breast carcinoma tissue.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	pH: 7.2 Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 40% Glycerol (glycerin, glycerine), 9.85% Tris glycine, 50% Tissue culture supernatant
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR3773

同种型IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab133248于以下的经测试应用
“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
WB	★★★★★ (1)	1/1000 - 1/10000. Predicted molecular weight: 23 kDa.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

应用说明Is unsuitable for Flow Cyt or IP.

靶标

功能Regulates the GDP/GTP exchange reaction of the Rho proteins by inhibiting the dissociation of GDP from them, and the subsequent binding of GTP to them.
序列相似性Belongs to the Rho GDI family.
细胞定位Cytoplasm.

图片



All lanes : Anti-RhoGDI antibody [EPR3773] (ab133248) at 1/1000 dilution

Lane 1 : Wild-type HEK293T cell lysate
Lane 2 : ARHGDIa knockout HEK293T cell lysate


Lysates/proteins at 20 µg per lane.
Performed under reducing conditions.

Predicted band size: 23 kDa

ab133248 was shown to react with aRHGDIA in wild-type HEK293T cells in Western blot with loss of signal observed in ARHGDIa knockout cell line **ab266447** (ARHGDIa knockout cell lysate **ab257356**). Wild-type HEK293T and ARHGDIa knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in

5% milk in TBST for 1 hr before incubation with ab133248 overnight at 4 °C at a 1/1000 dilution. Blots were incubated with goat anti-rabbit HRP secondary antibodies at 0.2µg/mL before imaging.

These data were provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.



Western blot - Anti-RhoGDI antibody [EPR3773] (ab133248)

All lanes : Anti-RhoGDI antibody [EPR3773] (ab133248) at 1/1000 dilution

Lane 1 : Wild-type HEK293T cell lysate

Lane 2 : ARHGDI knockout HEK293T cell lysate

Lane 3 : Jurkat cell lysate

Lane 4 : HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution

Predicted band size: 23 kDa

Observed band size: 23 kDa

Lanes 1-4: Merged signal (red and green). Green - ab133248 observed at 23 kDa. Red - loading control [ab8245](#) observed at 36 kDa.


ab133248 Anti-RhoGDI antibody [EPR3773] was shown to specifically react with RhoGDI in wild-type HEK293T cells. Loss of signal was observed when knockout cell line [ab266446](#) (knockout cell lysate [ab257355](#)) was used. Wild-type and RhoGDI knockout samples were subjected to SDS-PAGE. ab133248 and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated at room temperature for 2.5 hours at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RhoGDI antibody [EPR3773] (ab133248)

Immunohistochemical analysis of paraffin embedded Human breast carcinoma tissue labelling RhoGDI with ab133248 at 1/100 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-RhoGDI antibody [EPR3773] (ab133248)

All lanes : Anti-RhoGDI antibody [EPR3773] (ab133248) at 1/1000 dilution

Lane 1 : Wild-type HEK293T cell lysate

Lane 2 : ARHGDI knockout HEK293T cell lysate

Lane 3 : Jurkat cell lysate

Lane 4 : HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution


Predicted band size: 23 kDa

Observed band size: 27 kDa

Lanes 1-4: Merged signal (red and green). Green - ab133248 observed at 27 kDa. Red - loading control [ab8245](#) observed at 36 kDa.

ab133248 Anti-RhoGDI antibody [EPR3773] was shown to specifically react with RhoGDI in wild-type HEK293T cells. Loss of signal was observed when knockout cell line [ab266447](#) (knockout cell lysate [ab257356](#)) was used. Wild-type and RhoGDI knockout samples were subjected to SDS-PAGE. ab133248 and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse

IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



All lanes : Anti-RhoGDI antibody [EPR3773] (ab133248) at 1/1000 dilution

Lane 1 : Jurkat cell lysate

Lane 2 : HeLa cell lysate

Lane 3 : NIH3T3 cell lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 23 kDa

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-RhoGDI antibody [EPR3773] (ab133248)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors