

Anti-RPA32/RPA2 antibody [9H8] ab2175

★★★★★ [11 Abreviews](#) [223 References](#) [5 图像](#)

概述

产品名称	Anti-RPA32/RPA2抗体[9H8]
描述	小鼠单克隆抗体[9H8] to RPA32/RPA2
宿主	Mouse
经测试应用	适用于: Flow Cyt, WB, IHC-P
种属反应性	与反应: Human
免疫原	Full length native protein (purified) corresponding to Human RPA32/RPA2. BALB/C mice were injected with purified human RPA protein
阳性对照	WB: TE 671, SK N BE and HeLa whole cell lysates. IHC-P: Human tonsil tissue. Flow Cyt: HeLa cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.3</p> <p>Preservative: 0.05% Sodium azide</p> <p>Constituent: 1% BSA</p>
纯度	IgG fraction
克隆	单克隆
克隆编号	9H8
骨髓瘤	unknown
同种型	IgG1


kappa

The Abpromise guarantee **Abpromise™承诺保证使用ab2175于以下的经测试应用**

应用	Ab评论	说明
Flow Cyt	★★★★★ (1)	Use 1µg for 10 ⁶ cells. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
WB	★★★★★ (2)	Use at an assay dependent concentration. Predicted molecular weight: 32 kDa.
IHC-P		1/50 - 1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.


功能	<p>Required for DNA recombination, repair and replication. The activity of RP-A is mediated by single-stranded DNA binding and protein interactions.</p> <p>Functions as component of the alternative replication protein A complex (aRPA). aRPA binds single-stranded DNA and probably plays a role in DNA repair; it does not support chromosomal DNA replication and cell cycle progression through S-phase. In vitro, aRPA cannot promote efficient priming by DNA polymerase alpha but supports DNA polymerase delta synthesis in the presence of PCNA and replication factor C (RFC), the dual incision/excision reaction of nucleotide excision repair and RAD51-dependent strand exchange.</p>
翻译后修饰	<p>Phosphorylated in a cell-cycle-dependent manner (from the S phase until mitosis).</p> <p>Phosphorylated by ATR upon DNA damage, which promotes its translocation to nuclear foci. Can be phosphorylated in vitro by PRKDC/DNA-PK in the presence of Ku and DNA, and by CDK1.</p>
细胞定位	<p>Nucleus. Nucleus > PML body. Also present in PML nuclear bodies. Redistributes to discrete nuclear foci upon DNA damage.</p>

2




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RPA32/RPA2 antibody [9H8] (ab2175)

Formalin-fixed, paraffin-embedded human tonsil tissue stained for RPA32/RPA2 using ab2175 at 1/50 dilution in immunohistochemical analysis. Antigen retrieval with citrate buffer pH 6.0



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RPA32/RPA2 antibody [9H8] (ab2175)

Immunohistochemical analysis of formalin-fixed, paraffin-embedded Human tonsil tissue, staining RPA32/RPA2 with ab2175. Staining was detected using DAB.



Western blot - Anti-RPA32/RPA2 antibody [9H8]
(ab2175)

All lanes : Anti-RPA32/RPA2 antibody [9H8] (ab2175) at 5 µg/ml

Lane 1 : TE 671 (Human Rhabdomyosarcoma) Whole Cell Lysate

Lane 2 : SK N BE (Human neuroblastoma) Whole Cell Lysate

Lane 3 : HeLa (Human epithelial carcinoma cell line) Nuclear Lysate


Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Predicted band size: 32 kDa

Observed band size: 32 kDa




Flow Cytometry - Anti-RPA32/RPA2 antibody [9H8]
(ab2175)

This image is courtesy of an anonymous Abreview

Anti-RPA32/RPA2 antibody [9H8] (ab2175) is used to measure DNA Double-Stranded Breaks resection in U2OS cells +/- Camptotecin (CPT).

ab2175 is used at 1/200 dilution in 1xPBS + 0.2% Triton X-100 for 1 hour at 25°C.

Secondary antibody: anti-mouse Molecular Probes Alexa Fluor® 488 Conjugated at 1/200 dilution.



Flow Cytometry - Anti-RPA32/RPA2 antibody [9H8]
(ab2175)

Overlay histogram showing HeLa cells stained with ab2175 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab2175, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] ([ab91353](#), 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors