

Anti-RUNX2 antibody [EPR14334] ab192256

重组 RabMAb

★★★★★ **8 Abreviews** **92 References** **11 图像**

概述

产品名称	Anti-RUNX2抗体[EPR14334]
描述	兔单克隆抗体[EPR14334] to RUNX2
宿主	Rabbit
经测试应用	适用于: IHC-P, ICC/IF, Flow Cyt (Intra), ChIC/CUT&RUN-seq
种属反应性	与反应: Mouse, Human 预测可用于: Rat 
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	Human osteosarcoma, Human tonsil and Mouse spleen tissues; Saos-2 and PC cells.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 59% PBS, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR14334
同种型	IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab192256于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (5)	1/1000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		1/1000. For unpurified use at 1/500.
Flow Cyt (Intra)		1/50.
ChIC/CUT&RUN-seq		Use at an assay dependent concentration.

靶标

功能

Transcription factor involved in osteoblastic differentiation and skeletal morphogenesis. Essential for the maturation of osteoblasts and both intramembranous and endochondral ossification. CBF binds to the core site, 5'-PYGPYGGT-3', of a number of enhancers and promoters, including murine leukemia virus, polyomavirus enhancer, T-cell receptor enhancers, osteocalcin, osteopontin, bone sialoprotein, alpha 1(I) collagen, LCK, IL-3 and GM-CSF promoters (By similarity). Inhibits MYST4-dependent transcriptional activation.

组织特异性

Specifically expressed in osteoblasts.

疾病相关

Defects in RUNX2 are the cause of cleidocranial dysplasia (CLCD) [MIM:119600]; also known as cleidocranial dysostosis (CCD). CLCD is an autosomal dominant skeletal disorder with high penetrance and variable expressivity. It is due to defective endochondral and intramembranous bone formation. Typical features include hypoplasia/aplasia of clavicles, patent fontanelles, wormian bones (additional cranial plates caused by abnormal ossification of the calvaria), supernumerary teeth, short stature, and other skeletal changes. In some cases defects in RUNX2 are exclusively associated with dental anomalies.

序列相似性

Contains 1 Runt domain.

结构域

A proline/serine/threonine rich region at the C-terminus is necessary for transcriptional activation of target genes and contains the phosphorylation sites.


翻译后修饰

Phosphorylated; probably by MAP kinases (MAPK) (By similarity). Isoform 3 is phosphorylated on Ser-340.

细胞定位

Nucleus.

图片



Western blot - Anti-RUNX2 antibody [EPR14334] (ab192256)

All lanes : Anti-RUNX2 antibody [EPR14334] (ab192256) at 1/1000 dilution

Lane 1 : Saos-2 cell lysate

Lane 2 : MC3T3-E1 undifferentiated cell lysate

Lane 3 : MC3T3-E1 7-day Osteogenic differentiation cell lysate

Lane 4 : MC3T3-E1 14-day Osteogenic differentiation cell lysate

Lane 5 : MC3T3-E1 28-day Osteogenic differentiation cell lysate

Lane 6 : C2C12 cell lysate

Lane 7 : SH-SY5Y cell lysate


Lane 8 : NIH/3T3 cell lysate

Lane 9 : LNCaP cell lysate

Lysates/proteins at 20 µg per lane.

Observed band size: 60 kDa


Western blot: Anti-RUNX2 antibody [EPR14334] (ab192256) staining at 1/1000 dilution, shown in green; Mouse anti-CANX [CANX/1543] (**ab238078**) loading control staining at 1/20000 dilution, shown in magenta. In Western blot, ab192256 was shown to bind specifically to RUNX2. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution



ChIP/CUT&RUN sequencing - Anti-RUNX2 antibody [EPR14334] (ab192256)

ChIP/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/ μ L, 2.5×10^5 Saos-2 cells and 5 μ g of ab192256 [EPR14334]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.


The University of Geneva owns patents relevant to ChIP (Chromatin Immuno-Cleavage) methods.



ChIP/CUT&RUN sequencing - Anti-RUNX2 antibody [EPR14334] (ab192256)

ChIP/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/ μ L, 2.5×10^5 Saos-2 cells and 5 μ g of ab192256 [EPR14334]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.


The University of Geneva owns patents relevant to ChIP (Chromatin Immuno-Cleavage) methods.



ChIC/CUT&RUN sequencing - Anti-RUNX2 antibody
[EPR14334] (ab192256)

ChIC/CUT&RUN was performed using a pAG-MNase at a final concentration of 700 ng/ μ L, 2.5×10^5 Saos-2 cells and 5 μ g of ab192256 [EPR14334]. The resulting DNA was sequenced on the Illumina NovaSeq 6000 to a depth of 10 million reads. The negative IgG control **ab172730** is also shown.


The University of Geneva owns patents relevant to ChIC (Chromatin Immuno-Cleavage) methods.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RUNX2 antibody
[EPR14334] (ab192256)


Immunohistochemical analysis of paraffin-embedded Mouse spleen tissue labeling RUNX2 with ab192256 at 1/1000 dilution. A ready to use HRP Polymer for Rabbit IgG was used as the secondary. Hematoxylin counterstain.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-RUNX2 antibody [EPR14334] (ab192256)


Immunocytochemistry/Immunofluorescence analysis of Saos-2 (Human osteosarcoma cell line) labeling RUNX2 with purified ab192256 at 1/1000 dilution. Cells were fixed with 4% PFA and permeabilized with 0.1% tritonX-100. **ab150077** Goat anti rabbit IgG (Alexa Fluor[®]488) at 1/1000 was used as the secondary antibody. Nuclei were counterstained with DAPI. PBS was used instead of the primary antibody as the negative control.



Immunohistochemical analysis of paraffin-embedded Human osteosarcoma tissue labeling RUNX2 with ab192256 at 1/1000 dilution. A ready to use HRP Polymer for Rabbit IgG was used as the secondary. Hematoxylin counterstain.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RUNX2 antibody [EPR14334] (ab192256)




ab192256 staining RUNX2 in PC-3 (human prostate adenocarcinoma) by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde, permeabilised with 90% methanol and the sample was incubated with the primary antibody at a dilution of 1/50. A goat anti rabbit IgG (Alexa Fluor® 488) at a dilution of 1/2000 was used as the secondary antibody.

Isotype control: Rabbit monoclonal IgG (Black)

Unlabelled control: Cell without incubation with primary antibody and secondary antibody (Blue)


Flow Cytometry (Intracellular) - Anti-RUNX2 antibody [EPR14334] (ab192256)



Immunohistochemical analysis of paraffin-embedded Human tonsil tissue labeling RUNX2 with ab192256 at 1/1000 dilution. A ready to use HRP Polymer for Rabbit IgG was used as the secondary. Hematoxylin counterstain.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.





Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RUNX2 antibody [EPR14334] (ab192256)



Immunocytochemistry/ Immunofluorescence - Anti-RUNX2 antibody [EPR14334] (ab192256)

Immunofluorescent analysis of 4% formaldehyde fixed PC3 cells labeling RUNX2 using ab192256 at a 1/500 dilution. A Goat anti rabbit IgG (Alexa Fluor®488) **ab150077** was used as the secondary at a 1/200 dilution. Counterstain DAPI. Permeabilized using 0.1% Triton X-100. The two negative controls: 1. Primary ab concentration (anti-RUNX2) is 1:500 dilution, Secondary ab (Goat anti mouse IgG (Alexa Fluor®594)) is 1:500 dilution; 2. Primary ab concentration (anti-RUNX2) is 1:500 dilution, Secondary ab (Goat anti mouse IgG (Alexa Fluor®594)) is 1:500 dilution.

Why choose a recombinant antibody?

 Research with confidence Consistent and reproducible results	 Long-term and scalable supply Recombinant technology
 Success from the first experiment Confirmed specificity	 Ethical standards compliant Animal-free production

Anti-RUNX2 antibody [EPR14334] (ab192256)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors