


### Anti-S6K1 (phospho T389 + T412) antibody ab60948

★☆☆☆☆ [1 Abreviews](#) [16 References](#) [2 图像](#)

#### 概述

产品名称	Anti-S6K1 (phospho T389 + T412)抗体
描述	兔多克隆抗体to S6K1 (phospho T389 + T412)
宿主	Rabbit
特异性	ab60948 will detect both phospho-Thr389 of isoform alpha II and phospho-Thr412 of isoform alpha I.
经测试应用	适用于: ELISA, ICC/IF, IHC-P, WB
种属反应性	与反应: Human 预测可用于: Mouse, Rat 
免疫原	Synthetic peptide corresponding to Human S6K1 (phospho T389). Synthetic phosphopeptide derived from isoform 2 of human p70 S6 Kinase beta-1 (P23443) around the phosphorylation site of threonine 389 (G-F-TP-Y-V). Database link: <a href="#">P23443</a>
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine), 0.87% Sodium chloride  Without Mg <sup>2+</sup> and Ca <sup>2+</sup>
纯度	Immunogen affinity purified
纯化说明	ab60948 was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-

specific phosphopeptide. The antibody against non-phosphopeptide was removed by chromatography using non-phosphopeptide corresponding to the phosphorylation site.

**克隆** 多克隆  
**同种型** IgG

## 应用

**The Abpromise guarantee** **Abpromise™** 承诺保证使用ab60948于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ELISA		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
IHC-P		1/50 - 1/100.
WB		1/500 - 1/1000.

## 靶标

**功能** Acts to integrate nutrient and growth factor signals in regulation of protein synthesis, cell proliferation, cell growth, cell cycle progression and cell survival. Downstream effector of the mTOR signaling pathway. Phosphorylates specifically ribosomal protein S6 in response to insulin or several classes of mitogens. During translation initiation, the inactive form associates with the eIF-3 complex under conditions of nutrient depletion. Mitogenic stimulation leads to phosphorylation and dissociation from the eIF-3 complex and the free activated form can phosphorylate other translational targets including EIF4B. Promotes protein synthesis by phosphorylating PDCD4 at 'Ser-67' and targeting it for degradation. Phosphorylates RICTOR leading to regulation of mammalian target of rapamycin complex 2 (mTORC2) signaling; probably phosphorylates RICTOR at 'Thr-1135'. Phosphorylates IRS1 at multiple serine residues coupled with insulin resistance; probably phosphorylates IRS1 at 'Ser-270'. Required for TNF-alpha induced IRS-1 degradation. Phosphorylates EEF2K in response to IGF1 and inhibits EEF2K activity. Phosphorylates BAD at 'Ser-99' in response to IGF1 leading to BAD inactivation and inhibition of BAD-induced apoptosis. Phosphorylates mitochondrial RMP leading to dissociation of a RMP:PPP1CC complex; probably phosphorylates RMP at 'Ser-99'. The free mitochondrial PPP1CC can dephosphorylate RPS6KB1 at Thr-412 which is proposed to be a negative feed back mechanism for the RPS6KB1 antiapoptotic function. Phosphorylates GSK3B at 'Ser-9' under conditions leading to loss of the TSC1-TSC2 complex. Phosphorylates POLDIP3.


**组织特异性** Widely expressed.

**序列相似性** Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase subfamily.  
Contains 1 AGC-kinase C-terminal domain.  
Contains 1 protein kinase domain.

**结构域** The autoinhibitory domain is believed to block phosphorylation within the AGC-kinase C-terminal domain and the activation loop.  
The TOS (TOR signaling) motif is essential for activation by mTORC1.

翻译后修饰	Phosphorylation at Thr-412 is regulated by mTORC1. The phosphorylation at this site is maintained by an agonist-dependent autophosphorylation mechanism.
细胞定位	Cytoplasm; Nucleus. Cytoplasm and Cell junction > synapse > synaptosome. Mitochondrion outer membrane.

图片




Western blot - Anti-S6K1 (phospho T389 + T412) antibody (ab60948)

**All lanes :** Anti-S6K1 (phospho T389 + T412) antibody (ab60948)

**Lane 1 :** Jurkat cells treated with Insulin, 0.01U/ml

**Lane 2 :** Jurkat cells treated with Insulin, 0.01U/ml and phospho peptide used for blocking



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-S6K1 (phospho T389 + T412) antibody (ab60948)

ab60948 at 1/50 - 1/100 dilution staining P70 S6 Kinase beta in human breast carcinoma by Immunohistochemistry, Paraffin embedded tissue, in the absence or presence of the immunising peptide.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

**Our Abpromise to you: Quality guaranteed and expert technical support**

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors