

Anti-SERCA2 ATPase antibody [IID8] - BSA and Azide free ab255960

重组

4 图像

概述

产品名称	Anti-SERCA2 ATPase抗体[IID8] - BSA and Azide free
描述	小鼠单克隆抗体[IID8] to SERCA2 ATPase - BSA and Azide free
宿主	Mouse
特异性	Detects Sarcoplasmic or Endoplasmic Reticulum Calcium 2 (SERCA2) ATPase. This antibody does not discriminate between the two isoforms. By Western blot, this antibody detects an ~110 kDa protein representing SERCA2 ATPase from canine skeletal muscle triad preparations. Immunofluorescence staining of SECRA2 ATPase in rabbit skeletal muscle results in strong labeling of the entire type I (slow) myofiber consistent with sarcoplasmic reticulum localization. This antibody is not recommended for Western blot detection of rat SERCA2.
经测试应用	适用于: WB, IHC-P
种属反应性	与反应: Human
免疫原	Full length native protein (purified). This information is considered to be commercially sensitive.
阳性对照	IHC-P: Human skeletal muscle and cardiac muscle tissue. WB: Human skeletal muscle. HeLa, HepG2, A549 and A673 whole cell lysate.
常规说明	<p>ab255960 is the carrier-free version of ab2817.</p> <p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.2 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	IID8
同种型	IgG1

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab255960于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/2500. Detects a band of approximately 110 kDa (predicted molecular weight: 115 kDa).
IHC-P		Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

靶标

功能	This magnesium-dependent enzyme catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol to the sarcoplasmic reticulum lumen. Isoform 2 is involved in the regulation of the contraction/relaxation cycle.
组织特异性	Isoform 1 is widely expressed in smooth muscle and nonmuscle tissues such as in adult skin epidermis, with highest expression in liver, pancreas and lung, and intermediate expression in brain, kidney and placenta. Also expressed at lower levels in heart and skeletal muscle. Isoforms 2 and 3 are highly expressed in the heart and slow twitch skeletal muscle. Expression of isoform 3 is predominantly restricted to cardiomyocytes and in close proximity to the sarcolemma. Both isoforms are mildly expressed in lung, kidney, liver, pancreas and placenta. Expression of isoform 3 is amplified during monocytic differentiation and also observed in the fetal heart.
疾病相关	Defects in ATP2A2 are a cause of acrokeratosis verruciformis (AKV) [MIM:101900]; also known as Hopf disease. AKV is a localized disorder of keratinization, which is inherited as an autosomal dominant trait. Its onset is early in life with multiple flat-topped, flesh-colored papules on the hands and feet, punctate keratoses on the palms and soles, with varying degrees of nail involvement. The histopathology shows a distinctive pattern of epidermal features with hyperkeratosis, hypergranulosis, and acanthosis together with papillomatosis. These changes are frequently associated with circumscribed elevations of the epidermis that are said to resemble church

spires. There are no features of dyskeratosis or acantholysis, the typical findings in lesions of Darier disease.

Defects in ATP2A2 are the cause of Darier disease (DD) [MIM:124200]; also known as Darier-White disease (DAR). DD is an autosomal dominantly inherited skin disorder characterized by loss of adhesion between epidermal cells (acantholysis) and abnormal keratinization. Patients with mild disease may have no more than a few scattered keratotic papules or subtle nail changes, whereas those with severe disease are handicapped by widespread malodorous keratotic plaques. In a few families, neuropsychiatric abnormalities such as mild mental retardation, schizophrenia, bipolar disorder and epilepsy have been reported. Stress, UV exposure, heat, sweat, friction, and oral contraception exacerbate disease symptoms. Prevalence has been estimated at 1 in 50000. Clinical variants of DD include hypertrophic, vesicobullous, hypopigmented, cornifying, zosteriform or linear, acute and comedonal subtypes. Comedonal Darier disease (CDD) is characterized by the coexistence of acne-like comedonal lesions with typical Darier hyperkeratotic papules on light-exposed areas. At histopathologic level, CDD differs from classic DD in the prominent follicular involvement and the presence of greatly elongated dermal villi.

序列相似性

Belongs to the cation transport ATPase (P-type) (TC 3.A.3) family. Type IIA subfamily.


翻译后修饰

Nitrated under oxidative stress. Nitration on the two tyrosine residues inhibits catalytic activity.

细胞定位

Endoplasmic reticulum membrane. Sarcoplasmic reticulum membrane.

图片



All lanes : Anti-SERCA2 ATPase antibody [IID8] ([ab2817](#)) at 1/1000 dilution

Lane 1 : HeLa (human cervix adenocarcinoma epithelial cell), whole cell lysate

Lane 2 : HepG2 (human hepatocellular carcinoma epithelial cell), whole cell lysate

Lane 3 : A549 (human lung carcinoma epithelial cell), whole cell lysate

Lane 4 : A673 (human muscle Ewing's Sarcoma), whole cell lysate

Lane 5 : Human skeletal muscle

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Peroxidase-Conjugated Goat anti-Mouse IgG (H+L) at 1/100000 dilution

Predicted band size: 115 kDa


Observed band size: 110 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.

Exposure times: Lanes 1-2: 92 seconds; Lanes 3-4: 10 seconds; Lane 5: 3 seconds.

Samples are non-boiled as boiling may cause protein aggregates.

This data was developed using the same antibody clone in a different buffer formulation containing PBS and sodium azide ([ab2817](#)).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SERCA2 ATPase antibody [I1D8] - BSA and Azide free (ab255960)

Immunohistochemical analysis of paraffin-embedded human cardiac muscle tissue labeling SERCA2 ATPase with [ab2817](#) at 1/4000 dilution, followed by ready to use Goat Anti-Mouse IgG H&L (HRP polymer) ([ab214879](#)). Cytoplasmic staining on human cardiac muscle tissue is observed. Counterstained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, followed by ready to use Goat Anti-Mouse IgG H&L (HRP polymer) ([ab214879](#)).

Heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0).

This data was developed using the same antibody clone in a different buffer formulation containing PBS and sodium azide ([ab2817](#)).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SERCA2 ATPase antibody [I1D8] - BSA and Azide free (ab255960)

Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue labeling SERCA2 ATPase with [ab2817](#) at 1/4000 dilution, followed by ready to use Goat Anti-Mouse IgG H&L (HRP polymer) ([ab214879](#)). Cytoplasmic staining on human skeletal muscle tissue is observed. Counterstained with hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, followed by ready to use Goat Anti-Mouse IgG H&L (HRP polymer) ([ab214879](#)).

Heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0).

This data was developed using the same antibody clone in a different buffer formulation containing PBS and sodium azide ([ab2817](#)).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-SERCA2 ATPase antibody [IID8] - BSA and Azide free (ab255960)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors