

# Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] ab62322

**重组** RabMAb

**25 References** **6 图像**

## 概述

产品名称	Anti-SHP2 (phospho Y542)抗体[EP508(2)Y]
描述	兔单克隆抗体[EP508(2)Y] to SHP2 (phospho Y542)
宿主	Rabbit
经测试应用	<b>适用于:</b> ICC/IF, WB, IP <b>不适用于:</b> Flow Cyt
种属反应性	<b>与反应:</b> Mouse, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	Treated NIH/3T3 cell lysate.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> For more information <a href="#">see here</a> . Our RabMAb <sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a> . Rat: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EP508(2)Y

同种型	lgG
应用	
<p><b>The Abpromise guarantee</b>      <b>Abpromise™</b>承诺保证使用ab62322于以下的经测试应用</p> <p>“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。</p>	
应用	Ab评论
ICC/IF	说明
WB	1/100 - 1/250.
IP	1/1000. Detects a band of approximately 68 kDa (predicted molecular weight: 68 kDa). <b>For unpurified use at 1/50000- 1/100000.</b>
	1/30. <b>For unpurified use at 1/40</b>
应用说明	Is unsuitable for Flow Cyt.
靶标	
功能	Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus.
组织特异性	Widely expressed, with highest levels in heart, brain, and skeletal muscle.
疾病相关	<p>Defects in PTPN11 are the cause of LEOPARD syndrome type 1 (LEOPARD1) [MIM:151100]. It is an autosomal dominant disorder allelic with Noonan syndrome. The acronym LEOPARD stands for lentigines, electrocardiographic conduction abnormalities, ocular hypertelorism, pulmonic stenosis, abnormalities of genitalia, retardation of growth, and deafness.</p> <p>Defects in PTPN11 are the cause of Noonan syndrome type 1 (NS1) [MIM:163950]. Noonan syndrome (NS) is a disorder characterized by dysmorphic facial features, short stature, hypertelorism, cardiac anomalies, deafness, motor delay, and a bleeding diathesis. Some patients with Noonan syndrome type 1 develop multiple giant cell lesions of the jaw or other bony or soft tissues, which are classified as pigmented villomoduolar synovitis (PVNS) when occurring in the jaw or joints. Note=Mutations in PTPN11 account for more than 50% of the cases. Rarely, NS is associated with juvenile myelomonocytic leukemia (JMML). NS1 inheritance is autosomal dominant.</p> <p>Defects in PTPN11 are a cause of juvenile myelomonocytic leukemia (JMML) [MIM:607785]. JMML is a pediatric myelodysplastic syndrome that constitutes approximately 30% of childhood cases of myelodysplastic syndrome (MDS) and 2% of leukemia. It is characterized by leukocytosis with tissue infiltration and in vitro hypersensitivity of myeloid progenitors to granulocyte-macrophage colony stimulating factor.</p> <p>Defects in PTPN11 are a cause of metachondromatosis (MC) [MIM:156250]. It is a skeletal disorder with radiologic fetarures of both multiple exostoses and Ollier disease, characterized by the presence of multiple enchondromas and osteochondroma-like lesions.</p>
序列相似性	<p>Belongs to the protein-tyrosine phosphatase family. Non-receptor class 2 subfamily.</p> <p>Contains 2 SH2 domains.</p> <p>Contains 1 tyrosine-protein phosphatase domain.</p>
结构域	The SH2 domains repress phosphatase activity. Binding of these domains to phosphotyrosine-containing proteins relieves this auto-inhibition, possibly by inducing a conformational change in

## 翻译后修饰


the enzyme.

Phosphorylated on Tyr-546 and Tyr-584 upon receptor protein tyrosine kinase activation; which creates a binding site for GRB2 and other SH2-containing proteins.

## 细胞定位

Cytoplasm.

## 图片



ab62322 (purified) at 1:30 dilution (2µg) immunoprecipitating SHP2 in NIH/3T3 treated with 50ng/ml PDGF for 40min whole cell lysate.

Lane 1 (input): NIH/3T3 (Mouse embryonic fibroblast) treated with 50ng/ml PDGF for 40min whole cell lysate 10µg


Lane 2 (+): ab62322 & NIH/3T3 treated with 50ng/ml PDGF for 40min whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab62322 in NIH/3T3 treated with 50ng/ml PDGF for 40min whole cell lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP)

(**ab131366**) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDm/TBST.



**All lanes :** Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322) at 1/1000 dilution (Purified)

**Lane 1 :** NIH/3T3 (Mouse embryonic fibroblast) whole cell lysates

**Lane 2 :** NIH/3T3 (Mouse embryonic fibroblast) treated with 40ng/ml PDGF for 40 minutes whole cell lysates

**Lane 3 :** NIH/3T3 (Mouse embryonic fibroblast) treated with 40ng/ml PDGF for 40 minutes then incubated with phosphatase


Lysates/proteins at 15 µg per lane.

## Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

**Predicted band size:** 68 kDa


**Observed band size:** 68 kDa



Immunocytochemistry/ Immunofluorescence - Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322)

Ab62322 staining SHP2 in THP-1 cells ( Human monocytic leukaemia cell line) by ICC/IF (immunocytochemistry/immunofluorescence). Cells were fixed with 4% Paraformaldehyde and permeabilized with 0.1% TritonX-100. Samples were incubated with purified ab62322 at 8.8 µg/ml. Secondary antibody used was AlexaFluor®488 Goat anti-Rabbit (**ab150077**) at 2 µg/ml. Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594)( **ab195889**) used as counterstain at 2.5 µg/ml . DAPI was used as nuclear counterstain. Confocal image showing the expression was increased after treatment with TPA 200nM for 24h and PDGF 50ng/ml for 30min, the signal decreased after treatment with Lambda Protein Phosphatase 31I for 2h.

**This image was generated using the unpurified version of the product.**



Western blot - Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322)

**All lanes :** Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322) at 1/200 dilution

**Lane 1 :** THP-1 (Human monocytic leukemia monocyte) whole cell lysates

**Lane 2 :** THP-1 (Human monocytic leukemia monocyte) treated with Phorbol-12-myristate-13-acetate and platelet-derived growth factor.

**Lane 3 :** THP-1 (Human monocytic leukemia monocyte) treated with Phorbol-12-myristate-13-acetate and platelet-derived growth factor. Then the membrane was incubated with phosphatase.

Lysates/proteins at 15 µg per lane.

## Secondary

**All lanes :** Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution


**Predicted band size:** 68 kDa

**Observed band size:** 68 kDa

Blocking and diluting buffer used was 5% NFDM/TBST.

Purified batch of ab62322 was used.

**This image was generated using the unpurified version of the product.**



**All lanes :** Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322) at 1/50000 dilution

**Lane 1 :** NIH/3T3 cell lysates; untreated

**Lane 2 :** NIH/3T3 cell lysates; treated with PDGF

Lysates/proteins at 10 µg per lane.

#### Secondary

**All lanes :** goat anti-rabbit HRP conjugated, at 1/2000 dilution

**Predicted band size:** 68 kDa

**Observed band size:** 68 kDa

Beta Tubulin has been included as a loading control.

**This image was generated using the unpurified version of the product.**

Why choose a recombinant antibody?

<p><b>Research with confidence</b> Consistent and reproducible results</p>	<p><b>Long-term and scalable supply</b> Recombinant technology</p>
<p><b>Success from the first experiment</b> Confirmed specificity</p>	<p><b>Ethical standards compliant</b> Animal-free production</p>

Anti-SHP2 (phospho Y542) antibody [EP508(2)Y] (ab62322)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors