

Anti-Smad1 (phospho S463 + S465) antibody [EPR20662-20] ab226821

 **RabMAb**

4 References **5 图像**

概述

产品名称	Anti-Smad1 (phospho S463 + S465)抗体[EPR20662-20]
描述	兔单克隆抗体[EPR20662-20] to Smad1 (phospho S463 + S465)
宿主	Rabbit
特异性	Based on sequence homology this antibody also reacts with Smad5 (phospho S463/S465) and Smad9 (phospho S465/S467).
经测试应用	适用于: WB, Dot blot, ICC/IF, IP
种属反应性	与反应: Mouse, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HeLa grown in serum-free media overnight, treated with 100 ng/ml Calyculin A (ab141784) for 15min followed by Calyculin A removal and treatment with 100 ng/ml BMP2 for 30min, whole cell lysate; NIH/3T3 cultured in serum-free media overnight then treated with 50 ng/ml BMP2 for 30min whole cell lysate. ICC/IF: NIH3T3 cells FBS-deprived overnight before treatment with 50 ng/ml hBMP2 for 30min. IP: NIH/3T3 grown in serum-free media overnight then treated with 50 ng/ml BMP2 for 30min whole cell lysate.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.2

	Preservative: 0.01% Sodium azide
	Constituents: PBS, 0.05% BSA, 40% Glycerol (glycerin, glycerine)
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR20662-20
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab226821于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000. Detects a band of approximately 60 kDa (predicted molecular weight: 52 kDa).
Dot blot		1/1000.
ICC/IF		1/100.
IP		1/30.

靶标

功能	Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD1 is a receptor-regulated SMAD (R-SMAD). SMAD1/OAZ1/PSMB4 complex mediates the degradation of the CREBBP/EP300 repressor SNIP1.
组织特异性	Ubiquitous. Highest expression seen in the heart and skeletal muscle.
序列相似性	Belongs to the dwarfin/SMAD family. Contains 1 MH1 (MAD homology 1) domain. Contains 1 MH2 (MAD homology 2) domain.
翻译后修饰	Phosphorylated on serine by BMP type 1 receptor kinase. Ubiquitin-mediated proteolysis by SMAD-specific E3 ubiquitin ligase SMURF1.
细胞定位	Cytoplasm. Nucleus. Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4. Co-localizes with LEMD3 at the nucleus inner membrane.

图片



Western blot - Anti-Smad1 (phospho S463 + S465) antibody [EPR20662-20] (ab226821)

All lanes : Anti-Smad1 (phospho S463 + S465) antibody [EPR20662-20] (ab226821) at 1/1000 dilution

Lane 1 : HeLa (human epithelial cell line from cervix adenocarcinoma) grown in serum-free media overnight, whole cell lysate

Lane 2 : HeLa grown in serum-free media overnight, then treated with 100 ng/ml Calyculin A (**ab141784**) for 15 minutes, followed by Calyculin A removal and treatment with 100 ng/ml BMP2 for 30 minutes, whole cell lysate

Lane 3 : NIH/3T3 (mouse embryo fibroblast cell line) grown in serum-free media overnight, whole cell lysate

Lane 4 : NIH/3T3 cultured in serum-free media overnight, then treated with 50 ng/ml BMP2 for 30 minutes, whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution

Developed using the ECL technique.

Predicted band size: 52 kDa


Observed band size: 60 kDa

Exposure time:

Lanes 1 and 2: 3 minutes.

Lanes 3 and 4: 30 seconds.

Blocking/Dilution buffer: 5% NFDM/TBST.



Dot blot analysis of Smad1 (phospho S463 + S465) labeled with ab226821 at 1/1000 dilution.

Lane 1: Smad1 (phospho S463/S465) peptide;

Lane 2: Smad1 (phospho S463) peptide;

Lane 3: Smad1 (phospho S465) peptide;

Lane 4: Smad1 peptide (not phosphorylated);

Lane 5: Smad5 (phospho S463/S465) peptide;

Lane 6: Smad5 (phospho S463) peptide;

Lane 7: Smad5 (phospho S465) peptide;


Lane 8: Smad5 peptide (not phosphorylated).

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution was used as secondary antibody.

Blocking/Dilution buffer: 5% NFDM/TBST.

Exposure time: 3 minutes.


Based on sequence homology, this antibody cross-reacts with Smad5 (phospho S463/S465) and Smad9 (phospho S465/S467).



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (mouse embryo fibroblast cell line) cells labeling Smad1 (phospho S463 + S465) with ab226821 at 1/100 dilution, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) ([ab150077](#)) secondary antibody at 1/1000 dilution (green). Nuclear staining in hBMP2-treated NIH/3T3 cells. Cells were FBS-deprived overnight before treatment with 50 ng/ml hBMP2 for 30 minutes.

The nuclear counter stain is DAPI (blue). Tubulin is detected with Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) ([ab195889](#)) (red) at 1/200 dilution.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) ([ab150077](#)) secondary antibody at 1/1000 dilution.



Immunoprecipitation - Anti-Smad1 (phospho S463 + S465) antibody [EPR20662-20] (ab226821)

Smad 1 (phospho S463 + S465) was immunoprecipitated from 0.35 mg NIH/3T3 (mouse embryo fibroblast cell line) grown in serum-free media overnight, then treated with 50 ng/ml BMP2 for 30 minutes, whole cell lysate with ab226821 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab226821 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)), was used for detection at 1/1000 dilution.

Lane 1: NIH/3T3 grown in serum-free media overnight, then treated with 50 ng/ml BMP2 for 30 minutes, whole cell lysate 10 µg (Input).

Lane 2: ab226821 IP in NIH/3T3 grown in serum-free media overnight, then treated with 50 ng/ml BMP2 for 30 minutes, whole cell lysate.

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab226821 in NIH/3T3 grown in serum-free media overnight, then treated with 50 ng/ml BMP2 for 30 minutes, whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 10 seconds.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Smad1 (phospho S463 + S465) antibody
[EPR20662-20] (ab226821)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors