


# Anti-Smad4 antibody ab236321

[5 References](#) [5 图像](#)

### 概述

<b>产品名称</b>	Anti-Smad4抗体
<b>描述</b>	兔多克隆抗体to Smad4
<b>宿主</b>	Rabbit
<b>经测试应用</b>	<b>适用于:</b> IP, IHC-P, WB
<b>种属反应性</b>	<b>与反应:</b> Mouse, Human <b>预测可用于:</b> Rat, Cow, Pig 
<b>免疫原</b>	Recombinant fragment corresponding to Human Smad4 aa 1-200. Database link: <a href="#">Q13485</a> <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
<b>阳性对照</b>	WB: SH-SY5Y, 3T3 and HepG2 whole cell lysate. IHC-P: Human colon cancer and kidney tissue. IP: Jurkat cells.
<b>常规说明</b>	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 性能

<b>形式</b>	Liquid
<b>存放说明</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>存储溶液</b>	pH: 7.40 Preservative: 0.03% Proclin 300 Constituents: PBS, 50% Glycerol (glycerin, glycerine)
<b>纯度</b>	Protein G purified
<b>纯化说明</b>	Protein >95%.
<b>克隆</b>	多克隆

同种型

IgG

## 应用

### The Abpromise guarantee

**Abpromise™**承诺保证使用ab236321于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IP		1/200 - 1/2000.
IHC-P		1/20 - 1/200.
WB		1/500 - 1/2000. Predicted molecular weight: 60 kDa.

## 靶标

### 功能

Common SMAD (co-SMAD) is the coactivator and mediator of signal transduction by TGF-beta (transforming growth factor). Component of the heterotrimeric SMAD2/SMAD3-SMAD4 complex that forms in the nucleus and is required for the TGF-mediated signaling. Promotes binding of the SMAD2/SMAD4/FAST-1 complex to DNA and provides an activation function required for SMAD1 or SMAD2 to stimulate transcription. Component of the multimeric SMAD3/SMAD4/JUN/FOS complex which forms at the AP1 promoter site; required for synergistic transcriptional activity in response to TGF-beta. May act as a tumor suppressor.

### 疾病相关

Defects in SMAD4 are a cause of pancreatic cancer (PNCA) [MIM:260350].  
Defects in SMAD4 are a cause of juvenile polyposis syndrome (JPS) [MIM:174900]; also known as juvenile intestinal polyposis (JIP). JPS is an autosomal dominant gastrointestinal hamartomatous polyposis syndrome in which patients are at risk for developing gastrointestinal cancers. The lesions are typified by a smooth histological appearance, predominant stroma, cystic spaces and lack of a smooth muscle core. Multiple juvenile polyps usually occur in a number of Mendelian disorders. Sometimes, these polyps occur without associated features as in JPS; here, polyps tend to occur in the large bowel and are associated with an increased risk of colon and other gastrointestinal cancers.  
Defects in SMAD4 are a cause of juvenile polyposis/hereditary hemorrhagic telangiectasia syndrome (JP/HHT) [MIM:175050]. JP/HHT syndrome phenotype consists of the coexistence of juvenile polyposis (JIP) and hereditary hemorrhagic telangiectasia (HHT) [MIM:187300] in a single individual. JIP and HHT are autosomal dominant disorders with distinct and non-overlapping clinical features. The former, an inherited gastrointestinal malignancy predisposition, is caused by mutations in SMAD4 or BMPR1A, and the latter is a vascular malformation disorder caused by mutations in ENG or ACVRL1. All four genes encode proteins involved in the transforming-growth-factor-signaling pathway. Although there are reports of patients and families with phenotypes of both disorders combined, the genetic etiology of this association is unknown.  
Defects in SMAD4 may be a cause of colorectal cancer (CRC) [MIM:114500].

### 序列相似性

Belongs to the dwarfin/SMAD family.  
Contains 1 MH1 (MAD homology 1) domain.  
Contains 1 MH2 (MAD homology 2) domain.

### 结构域

The MH1 domain is required for DNA binding.  
The MH2 domain is required for both homomeric and heteromeric interactions and for

transcriptional regulation. Sufficient for nuclear import.


## 翻译后修饰

Monoubiquitinated on Lys-519 by E3 ubiquitin-protein ligase TRIM33. Monoubiquitination hampers its ability to form a stable complex with activated SMAD2/3 resulting in inhibition of TGF-beta/BMP signaling cascade. Deubiquitination by USP9X restores its competence to mediate TGF-beta signaling.

## 细胞定位

Cytoplasm. Nucleus. Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with R-SMAD.

## 图片



**All lanes :** Anti-Smad4 antibody (ab236321) at 1/500 dilution


**Lane 1 :** SH-SY5Y (human neuroblastoma cell line from bone marrow) whole cell lysate

**Lane 2 :** NIH/3T3 (mouse embryo fibroblast cell line) whole cell lysate


### Secondary

**All lanes :** Goat polyclonal to rabbit IgG at 1/10000 dilution

**Predicted band size:** 60 kDa




Paraffin-embedded human kidney tissue stained for Smad4 using ab236321 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Smad4 antibody (ab236321)

Paraffin-embedded human colon cancer tissue stained for Smad4 using ab236321 at 1/100 dilution in immunohistochemical analysis.




Western blot - Anti-Smad4 antibody (ab236321)

Anti-Smad4 antibody (ab236321) at 1/500 dilution + HepG2 (human liver hepatocellular carcinoma cell line) whole cell lysate

**Secondary**

Goat polyclonal to rabbit IgG at 1/10000 dilution

**Predicted band size:** 60 kDa



Smad4 was immunoprecipitated from 0.5 mg Jurkat (human T cell leukemia cell line from peripheral blood) whole cell lysate with ab236321.

**Lane 1:** Rabbit control IgG instead of ab236321 in Jurkat whole cell lysate.

**Lane 2:** ab236321 IP in Jurkat whole cell lysate.

**Lane 3:** Jurkat whole cell lysate 10 µg (Input).

For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody at 1/2000 dilution.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors