


Anti-SRC3 antibody ab2831

11 References **4 图像**

概述

产品名称	Anti-SRC3抗体
描述	兔多克隆抗体to SRC3
宿主	Rabbit
特异性	Detects amplified in breast cancer 1 (AIB1).
经测试应用	适用于: WB, ChIP, IHC-P, ICC/IF
种属反应性	与反应: Human 预测可用于: Xenopus laevis 
免疫原	Synthetic peptide corresponding to Human SRC3 aa 3-15. Sequence: GLGENLDPLASDS

(Peptide available as **ab4915**)

 **Run BLAST with**

 **Run BLAST with**

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituents: 0.1% BSA, 99% PBS
纯度	Immunogen affinity purified
克隆	多克隆

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab2831于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use a concentration of 2 µg/ml. Predicted molecular weight: 155 kDa. Can be blocked with AIB1 peptide or reuse.
ChIP		Use at an assay dependent concentration.
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF		Use at an assay dependent concentration.

靶标

功能	Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone-dependent fashion. Plays a central role in creating a multisubunit coactivator complex, which probably acts via remodeling of chromatin. Involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). Displays histone acetyltransferase activity. Also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit. Interacts with PSMB9.
组织特异性	Widely expressed. High expression in heart, skeletal muscle, pancreas and placenta. Low expression in brain, and very low in lung, liver and kidney.
序列相似性	Belongs to the SRC/p160 nuclear receptor coactivator family. Contains 1 basic helix-loop-helix (bHLH) domain. Contains 1 PAS (PER-ARNT-SIM) domain.
结构域	Contains three Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. Motifs 1 and 2 are essential for the association with nuclear receptors, and constitute the RID domain (Receptor-interacting domain).
翻译后修饰	Acetylated by CREBBP. Acetylation occurs in the RID domain, and disrupts the interaction with nuclear receptors and regulates its function. Methylated by CARM1. Phosphorylated by IKK complex. Regulated its function.
细胞定位	Cytoplasm. Nucleus. Mainly cytoplasmic and weakly nuclear. Upon TNF activation and subsequent phosphorylation, it translocates from the cytoplasm to the nucleus.

图片


Fig. 1

~155 kDa





Western blot - Anti-SRC3 antibody (ab2831)

Western blot of Human SRC3 in T47D cell lysate using ab2831.



Immunocytochemistry/ Immunofluorescence - Anti-SRC3 antibody (ab2831)

ICC/IF image of ab2831 stained human HeLa cells. The cells were methanol fixed (5 min) and incubated with the antibody (ab2831, 5µg/ml) for 1h at room temperature. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Image-iT™ FX Signal Enhancer was used as the primary blocking agent, 5% BSA (in TBS-T) was used for all other blocking steps. DAPI was used to stain the cell nuclei (blue). Alexa Fluor® 594 WGA was used to label plasma membranes (red).




ChIP - Anti-SRC3 antibody (ab2831)

This image is courtesy of Sylvain Daujat

ab2831 can be used in ChIP. You can expect recruitment of AIB1 on the pS2 promoter at some point during transcriptional activation. The amount of DNA precipitated is significant.

Sonicated Chromatin prepared from untreated or 17beta-estradiol (E2) treated MCF7 cells was subjected to the ChIP procedure with ab2831 to AIB1 and the immunoprecipitated chromatin was analysed in the proximal region of the estrogen-responsive pS2 promoter (as shown above) and quantified by real-time PCR (values are % of inputs). The primers are designed to follow the nucleosome E (including the Estrogen Responsive Element ERE). 3 µl of ab2831 and 2x10⁶ cells were used in each ChIP experiment.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SRC3 antibody (ab2831)

IHC image of ab2831 staining in human melanoma formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab2831, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors