

Anti-Streptococcus pyogenes Group A Carbohydrate antibody ab9191

10 References

概述

产品名称	抗化脓性链球菌Group A Carbohydrate抗体
描述	山羊多克隆抗体to化脓性链球菌Group A Carbohydrate
宿主	Goat
特异性	Streptococcus Pyogenes Group A
经测试应用	适用于: Dot blot, ELISA, ICC/IF, IHC-Fr, IP, WB, IHC-P
种属反应性	与反应: Streptococcus pyogenes
免疫原	Tissue, cells or virus corresponding to Streptococcus pyogenes Streptococcus pyogenes Group A Carbohydrate. Killed Streptococcus pyogenes vaccine Strain J17A4.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Constituent: 0.0268% PBS
纯度	Immunogen affinity purified
纯化说明	Serum was collected and antibody to the Group A carbohydrate was immunoaffinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee Abpromise™承诺保证使用ab9191于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Dot blot		Use at an assay dependent concentration.
ELISA		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. PubMed: 29416987

靶标

相关性 Streptococcus pyogenes (Group A streptococcus) is a Gram-positive, nonmotile, nonsporeforming coccus that occurs in chains or in pairs of cells. Individual cells are round-to-ovoid cocci, 0.6-1.0 micrometer in diameter. Streptococci divide in one plane and thus occur in pairs or (especially in liquid media or clinical material) in chains of varying lengths. The metabolism of S. pyogenes is fermentative; the organism is a catalase-negative aerotolerant anaerobe (facultative anaerobe), and requires enriched medium containing blood in order to grow. Group A streptococci typically have a capsule composed of hyaluronic acid and exhibit beta (clear) hemolysis on blood agar.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors