

Anti-Tau (phospho S404) antibody [EPR2605] ab92676

重组 RabMAb


★★★★★ **4 Abreviews** **29 References** **13 图像**

概述

产品名称	Anti-Tau (phospho S404)抗体[EPR2605]
描述	兔单克隆抗体[EPR2605] to Tau (phospho S404)
宿主	Rabbit
特异性	The specificity of this antibody refers to P10636-8.
经测试应用	适用于: IP, WB, IHC-P, IHC-Fr, ICC/IF, Dot blot
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	ICC/IF: Mouse primary cortical cultures; WB: Human, mouse and rat brain lysate; IHC: Mouse and rat cerebrum and Human glioma tissue; IP: Human brain lysate, IHC-Fr: Mouse cerebrum tissue.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at -20°C. Stable for 12 months at -20°C.
解离常数 (K _D)	K _D = 1.67 x 10 ⁻¹¹ M



存储溶液	pH: 7.20
------	----------

	Preservative: 0.05% Sodium azide
	Constituents: 40% Glycerol (glycerin, glycerine), 59% PBS, 0.01% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR2605
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab92676于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IP		1/20.
WB	★★★★★ (3)	1/500 - 1/2000. Predicted molecular weight: 79 kDa.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
IHC-Fr		1/50. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20)
ICC/IF	★★★★★ (1)	1/50.
Dot blot		Use at an assay dependent concentration.

靶标

功能	Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization.
组织特异性	Expressed in neurons. Isoform PNS-tau is expressed in the peripheral nervous system while the others are expressed in the central nervous system.
疾病相关	Note=In Alzheimer disease, the neuronal cytoskeleton in the brain is progressively disrupted and replaced by tangles of paired helical filaments (PHF) and straight filaments, mainly composed of hyperphosphorylated forms of TAU (PHF-TAU or AD P-TAU). Defects in MAPT are a cause of frontotemporal dementia (FTD) [MIM:600274]; also called frontotemporal dementia (FTD), pallido-ponto-nigral degeneration (PPND) or historically termed Pick complex. This form of frontotemporal dementia is characterized by presenile dementia with behavioral changes, deterioration of cognitive capacities and loss of memory. In some cases, parkinsonian symptoms are prominent. Neuropathological changes include frontotemporal atrophy often associated with atrophy of the basal ganglia, substantia nigra, amygdala. In most

cases, protein tau deposits are found in glial cells and/or neurons.

Defects in MAPT are a cause of Pick disease of the brain (PDB) [MIM:172700]. It is a rare form of dementia pathologically defined by severe atrophy, neuronal loss and gliosis. It is characterized by the occurrence of tau-positive inclusions, swollen neurons (Pick cells) and argentophilic neuronal inclusions known as Pick bodies that disproportionately affect the frontal and temporal cortical regions. Clinical features include aphasia, apraxia, confusion, anomia, memory loss and personality deterioration.

Note=Defects in MAPT are a cause of corticobasal degeneration (CBD). It is marked by extrapyramidal signs and apraxia and can be associated with memory loss. Neuropathologic features may overlap Alzheimer disease, progressive supranuclear palsy, and Parkinson disease.

Defects in MAPT are a cause of progressive supranuclear palsy type 1 (PSNP1) [MIM:601104, 260540]; also abbreviated as PSP and also known as Steele-Richardson-Olszewski syndrome. PSNP1 is characterized by akinetic-rigid syndrome, supranuclear gaze palsy, pyramidal tract dysfunction, pseudobulbar signs and cognitive capacities deterioration. Neurofibrillary tangles and gliosis but no amyloid plaques are found in diseased brains. Most cases appear to be sporadic, with a significant association with a common haplotype including the MAPT gene and the flanking regions. Familial cases show an autosomal dominant pattern of transmission with incomplete penetrance; genetic analysis of a few cases showed the occurrence of tau mutations, including a deletion of Asn-613.

序列相似性

Contains 4 Tau/MAP repeats.

发展阶段

Four-repeat (type II) tau is expressed in an adult-specific manner and is not found in fetal brain, whereas three-repeat (type I) tau is found in both adult and fetal brain.

结构域

The tau/MAP repeat binds to tubulin. Type I isoforms contain 3 repeats while type II isoforms contain 4 repeats.

翻译后修饰

Phosphorylation at serine and threonine residues in S-P or T-P motifs by proline-directed protein kinases (PDPK: CDK1, CDK5, GSK-3, MAPK) (only 2-3 sites per protein in interphase, seven-fold increase in mitosis, and in PHF-tau), and at serine residues in K-X-G-S motifs by MAP/microtubule affinity-regulating kinase (MARK) in Alzheimer diseased brains.

Phosphorylation decreases with age. Phosphorylation within tau's repeat domain or in flanking regions seems to reduce tau's interaction with, respectively, microtubules or plasma membrane components. Phosphorylation on Ser-610, Ser-622, Ser-641 and Ser-673 in several isoforms during mitosis.

Polyubiquitinated. Requires functional TRAF6 and may provoke SQSTM1-dependent degradation by the proteasome (By similarity). PHF-tau can be modified by three different forms of polyubiquitination. 'Lys-48'-linked polyubiquitination is the major form, 'Lys-6'-linked and 'Lys-11'-linked polyubiquitination also occur.

Glycation of PHF-tau, but not normal brain tau. Glycation is a non-enzymatic post-translational modification that involves a covalent linkage between a sugar and an amino group of a protein molecule forming ketoamine. Subsequent oxidation, fragmentation and/or cross-linking of ketoamine leads to the production of advanced glycation endproducts (AGES). Glycation may play a role in stabilizing PHF aggregation leading to tangle formation in AD.


细胞定位

Cytoplasm > cytosol. Cell membrane. Cytoplasm > cytoskeleton. Cell projection > axon. Mostly found in the axons of neurons, in the cytosol and in association with plasma membrane components.

形式

There are 9 isoforms produced by alternative splicing.

图片



Immunoprecipitation - Anti-Tau (phospho S404)
antibody [EPR2605] (ab92676)

Purified ab92676 at 1/20 dilution (0.5µg) immunoprecipitating Tau in Mouse brain lysate.

Lane 1 (input): Mouse brain lysate 10µg

Lane 2 (+): ab92676 + Mouse brain lysate.


Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab92676 in Mouse brain lysate.

VeriBlot for IP Detection Reagent (HRP) (**ab131366**) (1/1000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm/TBST.

Observed band size: 50-70 kDa



Immunoprecipitation - Anti-Tau (phospho S404)
antibody [EPR2605] (ab92676)

Purified ab92676 at 1/20 dilution (0.5µg) immunoprecipitating Tau in Human brain lysate.

Lane 1 (input): Human brain lysate 10µg

Lane 2 (+): ab92676 + Human brain lysate.


Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab92676 in Human brain lysate.

VeriBlot for IP Detection Reagent (HRP) (**ab131366**) (1/5000 dilution) was used for Western blotting.


Blocking Buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm/TBST.


Observed band size: 50-70 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human glioma tissue sections labeling Tau with purified ab92676 at 1/200 dilution (0.53 µg/mL). Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)

All lanes : Anti-Tau (phospho S404) antibody [EPR2605] (ab92676) at 1/1000 dilution (Purified)


Lane 1 : Rat brain lysate

Lane 2 : Rat brain lysate, the membrane treated with alkaline phosphatase for 1 hour

Secondary

All lanes : Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 79 kDa



Western blot - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)

All lanes : Anti-Tau (phospho S404) antibody [EPR2605] (ab92676) at 1/1000 dilution (Purified)


Lane 1 : Mouse brain lysate

Lane 2 : Mouse brain lysate, the membrane treated with alkaline phosphatase for 1 hour

Secondary

All lanes : Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 79 kDa



Western blot - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)

All lanes : Anti-Tau (phospho S404) antibody [EPR2605] (ab92676) at 1/1000 dilution (Purified)


Lane 1 : Human brain lysate

Lane 2 : Human brain lysate, the membrane treated with alkaline phosphatase for 1 hour

Secondary

All lanes : Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution


Predicted band size: 79 kDa



Immunocytochemistry/ Immunofluorescence - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)


Immunocytochemistry/ Immunofluorescence analysis of Embryonic mouse primary neural cells labeling Tau with purified ab92676 at 1:100 dilution (1 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) at 1:200 (2.5 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain.

Confocal image showing positive staining in neuron.




Immunohistochemistry (Frozen sections) - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)

Immunohistochemistry (Frozen sections) analysis of mouse cerebrum tissue sections labeling Tau with Purified ab92676 at 1/50 (1.9 µg/ml). Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. DAPI was used as a counterstain.



Immunocytochemistry/ Immunofluorescence - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)
This image is courtesy of an anonymous Abreview.

Mouse primary cortical cultures stained for Tau (phospho S404) (green) using ab92676 at 1/100 dilution in ICC/IF, followed by Donkey anti-Rabbit IgG (H+L) Alexa Fluor 488®.




Dot Blot - Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)

Dot blot analysis of Tau (pS404) phosphopeptide (Lane 1) and Tau non-phosphopeptide (Lane 2) labelling Tau (phospho S404) with ab92676 at a dilution of 1/1000. **ab97051** (Peroxidase conjugated goat anti-rabbit IgG (H+L)) was used as the secondary antibody at a dilution of 1/100000.

Blocking and dilution buffer: 5% NFDM/TBST.

Exposure time: 3 minutes.

Why choose a recombinant antibody?



Anti-Tau (phospho S404) antibody [EPR2605] (ab92676)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors