

Anti-TCR V beta 5 antibody [MEM-262] ab1898

3 References

概述

产品名称	Anti-TCR V beta 5抗体[MEM-262]
描述	小鼠单克隆抗体[MEM-262] to TCR V beta 5
宿主	Mouse
特异性	The antibody recognizes beta chains of the TCR expressed by HPB-ALL cell line [carrying V(beta5.3)] and a small subset of peripheral blood T cells. This subset is larger than that recognized by other V(beta5.3)-specific antibodies.
经测试应用	适用于: Flow Cyt, IP, WB
种属反应性	与反应: Human
免疫原	Tissue, cells or virus. This information is proprietary to Abcam and/or its suppliers.
常规说明	Functional Application Activates T cells (V beta 5-related subset). The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing. If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.097% Sodium azide Constituent: PBS
纯度	Protein A purified
纯化说明	Purified from hybridoma culture supernatant. Purity >95% by SDS-PAGE.
克隆	单克隆
克隆编号	MEM-262

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab1898于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration.

靶标

相关性 T cell receptors recognize foreign antigens which have been processed as small peptides and bound to major histocompatibility complex (MHC) molecules at the surface of antigen presenting cells (APC). Each T cell receptor is a dimer consisting of one alpha and one beta chain or one delta and one gamma chain. The beta locus includes V (variable), J (joining), diversity (D), and C (constant) segments. The Vβ5 TCR is a variant of the TCR β chain. It belongs to Ig superfamily and is expressed by a subset of peripheral blood T cells and some T cell clones. TCR Vβ5 has been reported to recognize some staphylococcus bacterial superantigens.

细胞定位 Cell Membrane

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors