

Anti-Thrombospondin 1 antibody [EPR22928-10] ab263905

重组 RabMAb

11 图像

概述

产品名称	Anti-Thrombospondin 1抗体[EPR22928-10]
描述	兔单克隆抗体[EPR22928-10] to Thrombospondin 1
宿主	Rabbit
经测试应用	适用于: WB, IHC-Fr, IP, IHC-P 不适用于: Flow Cyt (Intra) or ICC/IF
种属反应性	与反应: Mouse, Rat, Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: 3T3-L1 starved with 0.4% serum for 24 hours, then cultivated with 15% serum for 6 hours, whole cell lysate, Rat and mouse platelet whole cell lysate. IHC-P: Human spleen and bone marrow, Mouse and rat spleen tissue. IHC-Fr: Mouse and rat spleen tissue. IP: Mouse platelet cell lysate.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
纯度	Protein A purified
克隆	单克隆

克隆编号	EPR22928-10
同种型	IgG

应用

The Abpromise guarantee **Abpromise™承诺保证使用ab263905于以下的经测试应用**

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000. Detects a band of approximately 180 kDa (predicted molecular weight: 129 kDa).
IHC-Fr		1/500. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).
IP		1/30.
IHC-P		1/2000 - 1/5000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. Use at 1/5000 for human, 1/2000 for mouse and rat dilution.

应用说明 Is unsuitable for Flow Cyt (Intra) or ICC/IF.


靶标

功能	Adhesive glycoprotein that mediates cell-to-cell and cell-to-matrix interactions. Binds heparin. May play a role in dentinogenesis and/or maintenance of dentin and dental pulp (By similarity). Ligand for CD36 mediating antiangiogenic properties. Plays a role in ER stress response, via its interaction with the activating transcription factor 6 alpha (ATF6) which produces adaptive ER stress response factors.
----	---

序列相似性	<p>Belongs to the thrombospondin family.</p> <p>Contains 2 EGF-like domains.</p> <p>Contains 1 laminin G-like domain.</p> <p>Contains 1 TSP C-terminal (TSPC) domain.</p> <p>Contains 3 TSP type-1 domains.</p> <p>Contains 8 TSP type-3 repeats.</p> <p>Contains 1 VWFC domain.</p>
--------------	--

细胞定位 Endoplasmic reticulum. Sarcoplasmic reticulum.

图片




Immunohistochemistry (Frozen sections) - Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905)

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen rat spleen tissue labeling Thrombospondin 1 with ab263905 at 1/500 dilution followed by **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at a 1/1000 dilution. Positive staining on rat splenic megakaryocytes and platelet (PMID: 28239144) is observed. Nuclear counterstain DAPI.

Secondary antibody only control: Secondary antibody is **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at a 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).



Western blot - Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905)

All lanes : Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905) at 1/1000 dilution

Lane 1 : Mouse platelet whole cell lysate

Lane 2 : Rat platelet whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution

Predicted band size: 129 kDa

Observed band size: 180 kDa


Blocking buffer and concentration: 5% NFDm/TBST

Diluting buffer and concentration: 5% NFDm/TBST

Exposure time: Lane 1: 59 seconds Lane 2: 15 seconds

The identity of the lower MW bands at approximately 25 kDa is unknown.

The bands beneath the target band (~75-180 kDa) are likely to be degraded target fragments.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905)

Immunohistochemical analysis of paraffin-embedded rat spleen tissue labeling Thrombospondin 1 with ab263905 at 1/2000 dilution followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP Polymer). Positive staining on the megakaryocytes and platelets in the rat spleen. The section was incubated with ab263905 for 15 mins at room temperature. The immunostaining staining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP Polymer).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.



Immunoprecipitation - Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905)

Thrombospondin 1 was immunoprecipitated from 0.35 mg mouse platelet lysate 10µg with ab263905 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab263905 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used at 1/5000 dilution.


Lane 1: Mouse platelet lysate 10µg

Lane 2: ab263905 IP in mouse platelet lysate

Lane 3: Rabbit monoclonal IgG ([ab172730](#)) instead of ab263905 in mouse platelet lysate

Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 3 mins.




Immunohistochemistry (Frozen sections) - Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905)

Immunohistochemical analysis of 4% PFA-fixed, 0.2% Triton X-100 permeabilized frozen mouse spleen tissue labeling

Thrombospondin 1 with ab263905 at 1/500 dilution followed by **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at a 1/1000 dilution. Positive staining on mouse splenic megakaryocytes and platelet (PMID: 28239144) is observed. Nuclear counterstain DAPI.

Secondary antibody only control: Secondary antibody is **ab150077** AlexaFluor®488 Goat anti-Rabbit secondary at a 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905)

Immunohistochemical analysis of paraffin-embedded mouse spleen tissue labeling Thrombospondin 1 with ab263905 at 1/2000 dilution followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP Polymer). Positive staining on the megakaryocytes and platelets in the mouse spleen. The section was incubated with ab263905 for 15 mins at room temperature. The immunostaining staining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP Polymer).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905)

Immunohistochemical analysis of paraffin-embedded human spleen tissue labeling Thrombospondin 1 with ab263905 at 1/5000 dilution followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP Polymer). Positive staining on the platelets in the human spleen (PMID: 28239144). The section was incubated with ab263905 for 15 mins at room temperature. The immunostaining staining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP Polymer).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905)

Immunohistochemical analysis of paraffin-embedded human bone marrow tissue labeling Thrombospondin 1 with ab263905 at 1/5000 dilution followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP Polymer). Positive staining on the megakaryocytes in the human bone marrow (PMID: 28239144). The section was incubated with ab263905 for 15 mins at room temperature. The immunostaining staining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP Polymer).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 20 mins.



Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905) at 1/1000 dilution + Mouse platelet lysate at 20 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/25000 dilution


Predicted band size: 129 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.

The molecular weight observed is consistent with what has been described in the literature (PMID: 27588705).

Exposure time: 1 second.

Observed MW: 180 kDa.



All lanes : Anti-Thrombospondin 1 antibody [EPR22928-10] (ab263905) at 1/1000 dilution

Lane 1 : 3T3-L1 (mouse embryonic fibroblast) starved with 0.4% serum for 30 hours whole cell lysate

Lane 2 : 3T3-L1 starved with 0.4% serum for 24 hours, then cultivated with 15% serum for 6 hours. whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/25000 dilution

Predicted band size: 129 kDa

Observed band size: 180 kDa

Blocking and diluting buffer and concentration: 5% NFDm/TBST.

TSP 1 is a serum-responsive gene. Its express is elevated in quiescent 3T3 with serum addition.

The molecular weight observed is consistent with what has been described in the literature (PMID: 8422970, 27588705).

Exposure time: 3 minutes.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Thrombospondin 1 antibody [EPR22928-10]
(ab263905)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors