

### Anti-TSG101 antibody [EPR7130(B)] ab125011

重组 RabMAb


★★★★★ 10 Abreviews 426 References 10 图像

#### 概述

产品名称	Anti-TSG101抗体[EPR7130(B)]
描述	兔单克隆抗体[EPR7130(B)] to TSG101
宿主	Rabbit
经测试应用	适用于: WB, Flow Cyt (Intra), ICC/IF, IHC-P
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide within Human TSG101 aa 1-100. The exact sequence is proprietary.
阳性对照	WB: Mouse heart tissue lysates. K-562, Jurkat, SH-SY5Y, NIH/3T3 and PC-12 whole cell lysate. Mouse and rat heart and brain lysate. Flow Cyt: HeLa cells. ICC/IF: Jurkat cells and HeLa cells. IHC-P: Rat kidney tissue, mouse kidney tissue, human thyroid cancer tissue, human kidney tissue.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
解离常数 (K <sub>D</sub> )	K <sub>D</sub> = 4.00 x 10 <sup>-12</sup> M



[Learn more about K<sub>D</sub>](#)

存储溶液	pH: 7.20
------	----------

	Preservative: 0.01% Sodium azide
	Constituents: 40% Glycerol, 0.05% BSA, 59% PBS
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR7130(B)
同种型	IgG

应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab125011于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (7)	1/1000 - 1/10000. Detects a band of approximately 47, 52 kDa (predicted molecular weight: 44 kDa).
Flow Cyt (Intra)		1/20. For unpurified use at 1/3. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF	★★★★★ (3)	Use a concentration of 1 µg/ml. <b>For unpurified use at 1/10.</b> <b>This product gave a positive signal in HeLa fixed with 100% methanol (5 min).</b>
IHC-P		1/300. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See <b><u>IHC antigen retrieval protocols</u></b> . <b>For unpurified use at 1/30.</b>

靶标

功能	Component of the ESCRT-I complex, a regulator of vesicular trafficking process. Binds to ubiquitinated cargo proteins and is required for the sorting of endocytic ubiquitinated cargos into multivesicular bodies (MVBs). Mediates the association between the ESCRT-0 and ESCRT-I complex. Required for completion of cytokinesis; the function requires CEP55. May be involved in cell growth and differentiation. Acts as a negative growth regulator. Involved in the budding of many viruses through an interaction with viral proteins that contain a late-budding motif P-[ST]-A-P. This interaction is essential for viral particle budding of numerous retroviruses.
组织特异性	Heart, brain, placenta, lung, liver, skeletal, kidney and pancreas.
序列相似性	Belongs to the ubiquitin-conjugating enzyme family. UEV subfamily. Contains 1 SB (steadiness box) domain. Contains 1 UEV (ubiquitin E2 variant) domain.
结构域	The UEV domain is required for the interaction of the complex with ubiquitin. It also mediates the interaction with PTAP/PSAP motifs of HIV-1 P6 protein and human spumaretrovirus Gag protein. The coiled coil domain may interact with stathmin.

## 翻译后修饰


The UEV domain binds ubiquitin and P-[ST]-A-P peptide motif independently.

Monoubiquitinated at multiple sites by LRSAM1 and by MGRN1. Ubiquitination inactivates it, possibly by regulating its shuttling between an active membrane-bound protein and an inactive soluble form. Ubiquitination by MGRN1 requires the presence of UBE2D1.

## 细胞定位


Cytoplasm. Membrane. Nucleus. Late endosome membrane. Mainly cytoplasmic. Membrane-associated when active and soluble when inactive. Depending on the stage of the cell cycle, detected in the nucleus. Colocalized with CEP55 in the midbody during cytokinesis.

## 图片




Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labelling TSG101 with Purified ab125011 at 1:20 dilution (10 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1:2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabelled control - Cell without incubation with primary antibody and secondary antibody (Blue).

Flow Cytometry (Intracellular) - Anti-TSG101 antibody [EPR7130(B)] (ab125011)




Immunocytochemistry/ Immunofluorescence - Anti-TSG101 antibody [EPR7130(B)] (ab125011)

ab125011 staining TSG101 in HeLa cells. The cells were fixed with 100% methanol (5 min), permeabilised with 0.1% Triton x-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab125011 at 1 µg/ml and [ab7291](#), Mouse monoclonal [DM1A] to alpha Tubulin at 0.5 µg/ml. Cells were then incubated with [ab150081](#), Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and [ab150119](#), Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 647), pre-adsorbed at 1/1000 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue). Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.




Immunocytochemistry/ Immunofluorescence - Anti-TSG101 antibody [EPR7130(B)] (ab125011)

Immunocytochemistry analysis of Jurkat (Human T cell leukemia T lymphocyte) cells labeling TSG101 with Purified ab125011 at 1:50 dilution (4.5 µg/ml). Cells were fixed in 100% Methanol and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TSG101 antibody [EPR7130(B)] (ab125011)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat kidney tissue sections labeling TSG101 with Purified ab125011 at 1:300 dilution (0.73 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TSG101 antibody [EPR7130(B)] (ab125011)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse kidney tissue sections labeling TSG101 with Purified ab125011 at 1:300 dilution (0.73 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TSG101 antibody [EPR7130(B)] (ab125011)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human thyroid cancer tissue sections labeling TSG101 with Purified ab125011 at 1:300 dilution (0.73 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0) . Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TSG101 antibody [EPR7130(B)] (ab125011)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human kidney tissue sections labeling TSG101 with Purified ab125011 at 1:300 dilution (0.73 µg/ml). Heat mediated antigen retrieval was performed using Heat mediated antigen retrieval using Bond™ Epitope Retrieval Solution 2 (pH 9.0). Tissue was counterstained with Hematoxylin. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) secondary antibody was used. PBS instead of the primary antibody was used as the negative control.



Western blot - Anti-TSG101 antibody [EPR7130(B)] (ab125011)

**All lanes** : Anti-TSG101 antibody [EPR7130(B)] (ab125011) at 1/1000 dilution

**Lane 1** : K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate

**Lane 2** : Jurkat (Human T cell leukemia T lymphocyte) whole cell lysate

**Lane 3** : SH-SY5Y (Human neuroblastoma epithelial cell) whole cell lysate

**Lane 4** : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysate

**Lane 5** : PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysate

**Lane 6** : Mouse heart lysate

**Lane 7** : Rat heart lysate

**Lane 8** : Mouse brain lysate

**Lane 9** : Rat brain lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes** : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution


**Predicted band size:** 44 kDa

**Observed band size:** 47, 52 kDa

Blocking and dilution buffer: 5% NFDM/TBST.

The doublet are consistent with what have been described in the literature (PMID: 27882925, 31684046, 32932791, 11427703).

Image produced using the purified version.







Western blot - Anti-TSG101 antibody [EPR7130(B)]  
(ab125011)

Different batches of ab125011 were tested on Mouse heart lysate at 1.0 µg/ml. 15 µg of lysate was loaded in each lane. Bands observed at 45 kDa.

This data was generated using the unpurified format.

Why choose a recombinant antibody?

 <p><b>Research with confidence</b> Consistent and reproducible results</p>	 <p><b>Long-term and scalable supply</b> Recombinant technology</p>
 <p><b>Success from the first experiment</b> Confirmed specificity</p>	 <p><b>Ethical standards compliant</b> Animal-free production</p>

Anti-TSG101 antibody [EPR7130(B)] (ab125011)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors