


Anti-TSLP antibody ab115700

7 References **3 图像**

概述

产品名称	Anti-TSLP抗体
描述	兔多克隆抗体to TSLP
宿主	Rabbit
经测试应用	适用于: ELISA, WB, IHC-P
种属反应性	与反应: Mouse 预测可用于: Rat 
免疫原	Synthetic peptide corresponding to Mouse TSLP aa 70-120 (internal sequence) conjugated to keyhole limpet haemocyanin.
阳性对照	Mouse Liver and Spleen tissues; A20 cell lysate.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	<p>pH: 7.4</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituent: 99% PBS</p>
纯度	Protein A purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab115700于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ELISA		Use at an assay dependent concentration.
WB		Use a concentration of 0.5 - 2 µg/ml. Predicted molecular weight: 18 kDa.
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

靶标


功能	Cytokine that induces the release of T cell-attracting chemokines from monocytes and, in particular, enhances the maturation of CD11c(+) dendritic cells. Can induce allergic inflammation by directly activating mast cells.
组织特异性	Expressed in a number of tissues including heart, liver and prostate.
细胞定位	Secreted.

图片




ab115700, at 5 µg/ml, staining TSLP in Formalin-fixed, Paraffin-embedded Mouse Liver tissue by Immunohistochemistry followed by biotinylated secondary antibody, alkaline phosphatase-streptavidin and chromogen.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TSLP antibody (ab115700)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-TSLP antibody (ab115700)

ab115700, at 5 µg/ml, staining TSLP in Formalin-fixed, Paraffin-embedded Mouse Spleen tissue by Immunohistochemistry followed by biotinylated secondary antibody, alkaline phosphatase-streptavidin and chromogen.



Western blot - Anti-TSLP antibody (ab115700)

Lane 1 : Anti-TSLP antibody (ab115700) at 0.5 µg/ml

Lane 2 : Anti-TSLP antibody (ab115700) at 1 µg/ml

Lane 3 : Anti-TSLP antibody (ab115700) at 2 µg/ml

All lanes : A20 cell lysate

Predicted band size: 18 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors