

Anti-uPA Receptor/U-PAR antibody [EPR20191] ab218106

重组 RabMAb

4 References 6 图像

概述

产品名称	Anti-uPA Receptor/U-PAR抗体[EPR20191]
描述	兔单克隆抗体[EPR20191] to uPA Receptor/U-PAR
宿主	Rabbit
经测试应用	适用于: WB, IHC-P
种属反应性	与反应: Human
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: U937 whole cell lysate treated with 200nM PMA (Phorbol-12-myristate-13-acetate) for 72 hours. IHC-P: Human ovarian cancer, lung cancer and artery tissues.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.2</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR20191
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab218106于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000. Detects a band of approximately 35-65 kDa (predicted molecular weight: 37 kDa).
IHC-P		1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.


靶标

功能	Acts as a receptor for urokinase plasminogen activator. Plays a role in localizing and promoting plasmin formation. Mediates the proteolysis-independent signal transduction activation effects of U-PA. It is subject to negative-feedback regulation by U-PA which cleaves it into an inactive form.
组织特异性	Expressed in neurons of the rolandic area of the brain (at protein level). Expressed in the brain.
序列相似性	Contains 3 UPAR/Ly6 domains.
细胞定位	Secreted and Cell membrane.

图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-uPA Receptor/U-PAR antibody [EPR20191] (ab218106)



All lanes : Anti-uPA Receptor/U-PAR antibody [EPR20191] (ab218106) at 1/1000 dilution

Lane 1 : Untreated U937 (Human histiocytic lymphoma cell line) whole cell lysate

Lane 2 : U937 whole cell lysate treated with 200nM PMA (Phorbol-12-myristate-13-acetate) for 72 hours

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/100000 dilution


Predicted band size: 37 kDa

Observed band size: 35-65 kDa

Exposure time: 15 seconds

Blocking/Dilution buffer: 5% NFDM/TBST.

PMA-stimulated U937 cells express uPA Receptor/U-PAR in the range of 35kDa-60kDa. Non-stimulated U937 cells are negative. The deglycosylated form of the protein is at 35 kDa and the highly glycosylated protein is at 60kDa (PMID:24999729).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-uPA Receptor/U-PAR antibody [EPR20191] (ab218106)


Immunohistochemical analysis of paraffin-embedded human ovarian cancer tissue labeling uPA Receptor/U-PAR with ab218106 at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Membranous staining on the macrophages of human ovarian cancer is observed [PMID: 25575713].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-uPA Receptor/U-PAR antibody [EPR20191] (ab218106)


Immunohistochemical analysis of paraffin-embedded human lung cancer tissue labeling uPA Receptor/U-PAR with ab218106 at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Cytoplasmic staining on the stromal cells of human lung cancer is observed [PMID: 25575713].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-uPA Receptor/U-PAR antibody [EPR20191] (ab218106)

Immunohistochemical analysis of paraffin-embedded human artery tissue labeling uPA Receptor/U-PAR with ab218106 at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Positive staining on the neutrophils in human artery is observed [PMID: 25575713].

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-uPA Receptor/U-PAR antibody [EPR20191] (ab218106)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors