

Anti-Von Willebrand Factor antibody ab6994

★★★★★ [60 Abreviews](#) [457 References](#) [2 图像](#)

概述

产品名称	Anti-Von Willebrand Factor抗体
描述	兔多克隆抗体to Von Willebrand Factor
宿主	Rabbit
经测试应用	适用于: WB, ICC/IF, IHC-Fr, Flow Cyt, IHC-P, IHC-FoFr, IHC-FrFI
种属反应性	与反应: Rat, Sheep, Horse, Guinea pig, Cow, Dog, Human, Pig 不与反应: Chicken
免疫原	Full length native protein (purified) corresponding to Human Von Willebrand Factor. Purified from plasma.
阳性对照	IHC-P: Human kidney tissue.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.40 Preservative: 0.097% Sodium azide Constituent: PBS
纯度	IgG fraction
纯化说明	Whole antiserum is fractionated and then further purified by ion exchange chromatography to provide the IgG fraction of antiserum. This fraction is essentially free of other rabbit serum proteins.
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用 ab6994 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。

应用	Ab 评论	说明
WB	★★★★★ (10)	Use at an assay dependent concentration.
ICC/IF	★★★★★ (8)	Use at an assay dependent concentration.
IHC-Fr	★★★★★ (14)	Use at an assay dependent concentration.
Flow Cyt	★★★★★ (1)	Use at an assay dependent concentration. ab171870 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.
IHC-P	★★★★★ (22)	1/200 - 1/400. for IF and 1/1000-1/2000 for ABC methods with HRP conjugates. Perform enzymatic antigen retrieval with 0.1% pronase for 10 min at 35 °C before commencing with IHC protocol. Indirect Immunofluorescence: minimum working dilution of 1:200 was determined using FFPE sections of human tongue with FITC-conjugated secondary. Indirect Immunoperoxidase Labeling: minimum working dilution of 1:800 was determined
IHC-FoFr	★★★★★ (3)	Use at an assay dependent concentration. PubMed: 19622235
IHC-FrFI	★★★★★ (2)	Use at an assay dependent concentration. (see Abreview)


靶标

功能	Important in the maintenance of hemostasis, it promotes adhesion of platelets to the sites of vascular injury by forming a molecular bridge between sub-endothelial collagen matrix and platelet-surface receptor complex GPIb-IX-V. Also acts as a chaperone for coagulation factor VIII, delivering it to the site of injury, stabilizing its heterodimeric structure and protecting it from premature clearance from plasma.
组织特异性	Plasma.
疾病相关	Defects in VWF are the cause of von Willebrand disease (VWD) [MIM:277480]. VWD defines a group of hemorrhagic disorders in which the von Willebrand factor is either quantitatively or qualitatively abnormal resulting in altered platelet function. Symptoms vary depending on severity and disease type but may include prolonged bleeding time, deficiency of factor VIII and impaired platelet adhesion. Type I von Willebrand disease is the most common form and is characterized by partial quantitative plasmatic deficiency of an otherwise structurally and functionally normal Willebrand factor; type II is associated with a qualitative deficiency and functional anomalies of the Willebrand factor; type III is the most severe form and is characterized by total or near-total absence of Willebrand factor in the plasma and cellular compartments, also leading to a profound deficiency of plasmatic factor VIII.
序列相似性	Contains 1 CTCK (C-terminal cystine knot-like) domain.

Contains 4 TIL (trypsin inhibitory-like) domains.
Contains 3 VWFA domains.
Contains 3 VWFC domains.
Contains 4 VWFD domains.


结构域	The von Willebrand antigen 2 is required for multimerization of vWF and for its targeting to storage granules.
翻译后修饰	All cysteine residues are involved in intrachain or interchain disulfide bonds. N- and O-glycosylated.
细胞定位	Secreted. Secreted > extracellular space > extracellular matrix. Localized to storage granules.

图片



Immunohistochemical analysis of Formalin fixed paraffin-embedded sections human kidney tissue labeling Von Willebrand Factor with ab6994 at 1/2000.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Von Willebrand Factor antibody (ab6994)



Immunohistochemical analysis of Formalin fixed paraffin-embedded sections human kidney tissue labeling Von Willebrand Factor with ab6994 at 1/2000.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Von Willebrand Factor antibody (ab6994)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors