

Recombinant Human c-Maf protein ab114469

1 图像

描述		
产品名称	重组人c-Maf蛋白	
表达系统	Wheat germ	
Accession	<u>O75444-1</u>	
蛋白长度	Protein fragment	
无动物成分	No	
性质	Recombinant	
种属	Human	
序列	SCRFKRVQQRHVLESEKNQLLQQVDHLKQEISRLVRERDAYK EKYEKLVS SGFRENGSSSDNPSSPEFFITEPTRKLEPSVGYATFWKPQHR VLTSVFTK	
预测分子量	37 kDa including tags	
氨基酸	304 to 403	


技术指标		
Our <b>Abpromise guarantee</b> covers the use of <b>ab114469</b> in the following tested applications.		
The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.		
应用	SDS-PAGE  ELISA  Western blot	
形式	Liquid	

制备和贮存		
稳定性和存储	Shipped on dry ice. Upon delivery aliquot and store at -80°C. Avoid freeze / thaw cycles.  pH: 8.00 Constituents: 0.3% Glutathione, 0.79% Tris HCl	

常规信息		
------	--	--

功能	Acts as a transcriptional activator or repressor. Involved in embryonic lens fiber cell development. Recruits the transcriptional coactivators CREBBP and/or EP300 to crystallin promoters leading to up-regulation of crystallin gene during lens fiber cell differentiation. Activates the expression of IL4 in T helper 2 (Th2) cells. Increases T cell susceptibility to apoptosis by interacting with MYB and decreasing BCL2 expression. Together with PAX6, transactivates strongly the glucagon gene promoter through the G1 element. Activates transcription of the CD13 proximal promoter in endothelial cells. Represses transcription of the CD13 promoter in early stages of myelopoiesis by affecting the ETS1 and MYB cooperative interaction. Involved in the initial chondrocyte terminal differentiation and the disappearance of hypertrophic chondrocytes during endochondral bone development. Binds to the sequence 5'-[GT]G[GC]N[GT]NCTCAGNN-3' in the L7 promoter. Binds to the T-MARE (Maf response element) sites of lens-specific alpha- and beta-crystallin gene promoters. Binds element G1 on the glucagon promoter. Binds an AT-rich region adjacent to the TGC motif (atypical Maf response element) in the CD13 proximal promoter in endothelial cells (By similarity). When overexpressed, represses anti-oxidant response element (ARE)-mediated transcription. Involved either as an oncogene or as a tumor suppressor, depending on the cell context. Binds to the ARE sites of detoxifying enzyme gene promoters.
组织特异性	Expressed in endothelial cells.
疾病相关	<p>Note=A chromosomal aberration involving MAF is found in some forms of multiple myeloma (MM). Translocation t(14;16)(q32.3;q23) with an IgH locus.</p> <p>Defects in MAF are the cause of cataract pulverulent juvenile-onset MAF-related (CAPJOM) [MIM:610202]. A form of cataract with nuclear or cortical pulverulent opacities. Pulverulent cataracts are characterized by a dust-like, 'pulverised' appearance of the opacities which can be found in any part of the lens. The phenotype shows significant intra- and interfamilial variation, both in the distribution of the cataract and the degree of opacification. Some patients with cataract pulverulent juvenile-onset can present microcornea and bilateral iris colobomas in addition to cataract.</p> <p>Defects in MAF are the cause of cataract congenital cerulean type 4 (CCA4) [MIM:610202]. A cerulean form of congenital cataract. Cerulean cataracts are characterized by peripheral bluish and white opacifications organized in concentric layers with occasional central lesions arranged radially. The opacities are observed in the superficial layers of the fetal nucleus as well as the adult nucleus of the lens. Involvement is usually bilateral. Visual acuity is only mildly reduced in childhood. In adulthood, the opacifications may progress, making lens extraction necessary. Histologically the lesions are described as fusiform cavities between lens fibers which contain a deeply staining granular material. Although the lesions may take on various colors, a dull blue is the most common appearance and is responsible for the designation cerulean cataract.</p>
序列相似性	<p>Belongs to the bZIP family. Maf subfamily.</p> <p>Contains 1 bZIP domain.</p>
翻译后修饰	<p>Ubiquitinated, leading to its degradation by the proteasome. Ubiquitination is triggered by glucocorticoids.</p> <p>Phosphorylated by GSK3 and MAPK13 on serine and threonine residues (Probable). The phosphorylation status can serve to either stimulate or inhibit transcription.</p>
细胞定位	Nucleus.

图片



SDS-PAGE - Recombinant Human c-Maf protein  
(ab114469)

ab114469 analysed on a 12.5% SDS-PAGE stained with Coomassie Blue.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors