

Recombinant human CD86 protein (Fc Chimera) ab167720

[1 References](#) [7 图像](#)

描述

产品名称	重组人CD86蛋白(Fc Chimera)
生物活性	<p>Measured by its ability to induce IL2 secretion by Jurkat Human acute T cell leukemia cells. The ED₅₀ for this effect is typically 0.04 - 0.2 ng/ml.</p> <p>Measured by its binding ability in a functional ELISA. Immobilized Biotinylated Human / Cynomolgus / Rhesus macaque CD28, His,Avitag at 1 µg/mL (100 µL/well) on Streptavidin precoated (0.5 µg/well) plate, can bind Human ab167720 with a linear range of 2-78 ng/mL.</p> <p>Measured by its binding ability in a functional ELISA. Immobilized Biotinylated Mouse CD28, His,Avitag at 1 µg/mL (100 µL/well) on Streptavidin precoated (0.5 µg/well) plate, can bind ab167720 with a linear range of 5-78 ng/mL.</p> <p>Measured by its binding ability in a functional ELISA. Immobilized ab167720 at 5 µg/mL (100 µL/well) can bind Biotinylated Human CTLA-4, Fc Tag, Avi Tag with a linear range of 0.6-10 ng/mL.</p>
纯度	> 95 % SDS-PAGE.
内毒素水平	< 1.000 Eu/µg
表达系统	HEK 293 cells
Accession	<u>P42081</u>
蛋白长度	Protein fragment
无动物成分	No
性质	Recombinant
种属	Human
预测分子量	52 kDa
氨基酸	26 to 247
标签	Fc tag C-Terminus
额外的序列信息	Protein carries a human IgG1 Fc tag at the C-terminus. Due to glycosylation, the protein migrates as 66-86 kDa under reducing condition, and 120-140 kDa under non-reducing condition (SDS-PAGE).

技术指标

Our **Abpromise guarantee** covers the use of **ab167720** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Functional Studies SDS-PAGE ELISA
形式	Lyophilized

制备和贮存

稳定性和存储 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.

pH: 7.4

Constituents: 0.75% Glycine, 0.48% Tris, L-Arginine, Sodium chloride

Lyophilized from 0.22 µm filtered solution.

5-10% trehalose is commonly used for freeze drying, and after reconstitution, the trehalose is mostly about 3-5%

This product is an active protein and may elicit a biological response in vivo, handle with caution.

复溶 It is recommended to reconstitute the lyophilized protein with 500 µL sterile deionized water to a stock solution of 400 µg/mL.

常规信息

功能 Receptor involved in the costimulatory signal essential for T-lymphocyte proliferation and interleukin-2 production, by binding CD28 or CTLA-4. May play a critical role in the early events of T-cell activation and costimulation of naive T-cells, such as deciding between immunity and anergy that is made by T-cells within 24 hours after activation. Isoform 2 interferes with the formation of CD86 clusters, and thus acts as a negative regulator of T-cell activation.

组织特异性 Expressed by activated B-lymphocytes and monocytes.


序列相似性 Contains 1 Ig-like C2-type (immunoglobulin-like) domain.

Contains 1 Ig-like V-type (immunoglobulin-like) domain.

翻译后修饰 Polyubiquitinated; which is promoted by MARCH8 and results in endocytosis and lysosomal degradation.


细胞定位 Membrane.

图片




Human B7-2, Fc Tag on SDS-PAGE under reducing (R) and non-reducing (NR) conditions. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%. The protein migrates as 66-86 kDa under reducing (R) condition, and 120-140 kDa under non-reducing (NR) condition (SDS-PAGE).

SDS-PAGE - Recombinant human CD86 protein (ab167720)




FACS assay shows that Human B7-2, Fc Tag can bind to 293T cells overexpressing human CTLA4 protein. The concentration of Human B7-2 is 3 µg/ml.

Functional Studies - Recombinant human CD86 protein (ab167720)




FACS analysis shows that the binding of Human B7-2, Fc Tag to CD28 expressed on Jurkat E6.1 was inhibited by increasing concentration of neutralizing anti-CD28 antibody. The concentration of B7-2 used is 2 µg/ml. The IC50 is 0.022 µg/ml.

Functional Studies - Recombinant human CD86 protein (ab167720)




Flow Cytometry assay shows that recombinant Human B7-2, Fc Tag can bind to CD28 expressed on Jurkat E6.1. The concentration of B7-2 used is 2 µg/ml(Routinely tested).

Functional Studies - Recombinant human CD86 protein (ab167720)




Immobilized Biotinylated Human / Cynomolgus / Rhesus macaque CD28, His,Avitag at 1 µg/mL (100 µL/well) on Streptavidin precoated (0.5 µg/well) plate, can bind Human ab167720 with a linear range of 2-78 ng/mL.

ELISA - Recombinant human CD86 protein (Fc Chimera) (ab167720)



Immobilized Biotinylated Mouse CD28, His,Avitag at 1 µg/mL (100 µL/well) on Streptavidin precoated (0.5 µg/well) plate, can bind ab167720 with a linear range of 5-78 ng/mL.

ELISA - Recombinant human CD86 protein (Fc Chimera) (ab167720)



Immobilized ab167720 at 5 µg/mL (100 µL/well) can bind Biotinylated Human CTLA-4, Fc Tag, Avi Tag with a linear range of 0.6-10 ng/mL.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors