

Recombinant human IGF1 Receptor protein ab155622

4 图像

描述	
产品名称	重组人IGF1 Receptor蛋白
生物活性	Measured by its ability to bind Human IGF-I in a functional ELISA.
纯度	> 95 % SDS-PAGE.
内毒素水平	< 1.000 Eu/μg
表达系统	HEK 293 cells
Accession	<u>P08069</u>
蛋白长度	Protein fragment
无动物成分	No
性质	Recombinant
种属	Human
序列	EICGPGIDIRNDYQQLKRLNCTVIEGYLHILLISKAEDYRS YRFPKLTV ITEYLLLFRVAGLESLGDLFPNLTVIRGWKLFYNYALVIFEM TNLKDIGL YNLRNITRGAIRIEKNADLCYLSTVDWSLILDAVSNNYIVGN KPPKECGD LCPGTMEЕКPMCEKTTINNEYNYRCWTTNRCQKMCPSTCGKR ACTENNEC CHPECLGSCSAPDNDTACVACRHHYYAGVCVPACPPNTYRFE GWRCVDRD FCANILSAESSDSEGFVIHDGECMQECPSGFIRNGSQSMYCI PCEGPCPK VCEEEKTKTIDSVTSAQMLQGCTIFKGNLLINIRRGNNIAS ELENFMGL IEVVTGYVKIRHSHALVSLSFLKNLRLILGEEQLEGNYSFYV LDNQNLQQ LWDWDHRNLTIKAGKMYFAFNPKLCVSEIYRMEEVTGTKGRQ SKGDINTR NNGERASCESDVLHFTSTTTSKNRIIITWHRYRPPDYRDLIS FTVYYKEA PFKNVTEYDGQDACGSNSWNMVDVDLPPNKDVEPGILLHGLK PWTQYAVY VKAVTLTMVENDHIRGAKSEILYIRTNASVPSIPLDVLSASN

SSSQLIVK
WNPPSLPNGNLSYYIVRWQRQPQDGYLYRHNYCSKDKIPRK
YADGTIDI
EEVTENPKTEVCGGEKGPCCACPKTEAEKQAEKEEAERYKVF
ENFLHNSI
FVPRPERKRRDVMQVANTTSSRSRNTTAADTYNITDPEELE
TEYPPFES
RVDNKERTVISNLRPFTLYRIDIHSCNHEAEKLGCSASNFVF
ARTMPAEG
ADDIPGPVTWEPRPENSIFLKWPEPENPNGLILMYEIKYGSQ
VEDQRECV
SRQEYRKYGGAKLNRLNPGNYTARIQATSLSGNGSWTDPVFF
YVQAKTGY EN

预测分子量	104 kDa including tags
氨基酸	31 to 932
标签	His tag C-Terminus

技术指标

Our **Abpromise guarantee** covers the use of **ab155622** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	Functional Studies ELISA SDS-PAGE
形式	Lyophilized
补充说明	Measured by its ability to bind Human IGF-I in a functional ELISA.

制备和贮存

稳定性和存储	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle. pH: 7.40 Constituents: 95% PBS, 5% Trehalose This product is an active protein and may elicit a biological response in vivo, handle with caution.
复溶	It is recommended to reconstitute the lyophilized protein in sterile deionized water to a final concentration of 400 ug/ml. Solubilize for 30 to 60 minutes at room temperature with occasional gentle mixing. Carrier protein (0.1% HSA or BSA) is strongly recommended for further dilution and long term storage.

常规信息

功能	Receptor tyrosine kinase which mediates actions of insulin-like growth factor 1 (IGF1). Binds IGF1 with high affinity and IGF2 and insulin (INS) with a lower affinity. The activated IGF1R is involved in cell growth and survival control. IGF1R is crucial for tumor transformation and survival of malignant cell. Ligand binding activates the receptor kinase, leading to receptor autophosphorylation, and tyrosines phosphorylation of multiple substrates, that function as signaling adapter proteins
----	---

including, the insulin-receptor substrates (IRS1/2), Shc and 14-3-3 proteins. Phosphorylation of IRSs proteins lead to the activation of two main signaling pathways: the PI3K-AKT/PKB pathway and the Ras-MAPK pathway. The result of activating the MAPK pathway is increased cellular proliferation, whereas activating the PI3K pathway inhibits apoptosis and stimulates protein synthesis. Phosphorylated IRS1 can activate the 85 kDa regulatory subunit of PI3K (PIK3R1), leading to activation of several downstream substrates, including protein AKT/PKB. AKT phosphorylation, in turn, enhances protein synthesis through mTOR activation and triggers the antiapoptotic effects of IGF1R through phosphorylation and inactivation of BAD. In parallel to PI3K-driven signaling, recruitment of Grb2/SOS by phosphorylated IRS1 or Shc leads to recruitment of Ras and activation of the ras-MAPK pathway. In addition to these two main signaling pathways IGF1R signals also through the Janus kinase/signal transducer and activator of transcription pathway (JAK/STAT). Phosphorylation of JAK proteins can lead to phosphorylation/activation of signal transducers and activators of transcription (STAT) proteins. In particular activation of STAT3, may be essential for the transforming activity of IGF1R. The JAK/STAT pathway activates gene transcription and may be responsible for the transforming activity. JNK kinases can also be activated by the IGF1R. IGF1 exerts inhibiting activities on JNK activation via phosphorylation and inhibition of MAP3K5/ASK1, which is able to directly associate with the IGF1R. When present in a hybrid receptor with INSR, binds IGF1. PubMed:12138094 shows that hybrid receptors composed of IGF1R and INSR isoform Long are activated with a high affinity by IGF1, with low affinity by IGF2 and not significantly activated by insulin, and that hybrid receptors composed of IGF1R and INSR isoform Short are activated by IGF1, IGF2 and insulin. In contrast, PubMed:16831875 shows that hybrid receptors composed of IGF1R and INSR isoform Long and hybrid receptors composed of IGF1R and INSR isoform Short have similar binding characteristics, both bind IGF1 and have a low affinity for insulin.

组织特异性

Found as a hybrid receptor with INSR in muscle, heart, kidney, adipose tissue, skeletal muscle, hepatoma, fibroblasts, spleen and placenta (at protein level). Expressed in a variety of tissues. Overexpressed in tumors, including melanomas, cancers of the colon, pancreas prostate and kidney.

疾病相关

Insulin-like growth factor 1 resistance

序列相似性


Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily. Contains 4 fibronectin type-III domains. Contains 1 protein kinase domain.

翻译后修饰


Autophosphorylated on tyrosine residues in response to ligand binding. Autophosphorylation occurs in trans, i.e. one subunit of the dimeric receptor phosphorylates tyrosine residues on the other subunit. Autophosphorylation occurs in a sequential manner; Tyr-1165 is predominantly phosphorylated first, followed by phosphorylation of Tyr-1161 and Tyr-1166. While every single phosphorylation increases kinase activity, all three tyrosine residues in the kinase activation loop (Tyr-1165, Tyr-1161 and Tyr-1166) have to be phosphorylated for optimal activity. Can be autophosphorylated at additional tyrosine residues (in vitro). Autophosphorylated is followed by phosphorylation of juxtamembrane tyrosines and C-terminal serines. Phosphorylation of Tyr-980 is required for IRS1- and SHC1-binding. Phosphorylation of Ser-1278 by GSK-3 β restrains kinase activity and promotes cell surface expression, it requires a priming phosphorylation at Ser-1282. Dephosphorylated by PTPN1. Polyubiquitinated at Lys-1168 and Lys-1171 through both 'Lys-48' and 'Lys-29' linkages, promoting receptor endocytosis and subsequent degradation by the proteasome. Ubiquitination is facilitated by pre-existing phosphorylation. Sumoylated with SUMO1. Controlled by regulated intramembrane proteolysis (RIP). Undergoes metalloprotease-dependent constitutive ectodomain shedding to produce a membrane-anchored 52 kDa C-Terminal fragment which is further processed by presenilin gamma-secretase to yield an intracellular 50

	kDa fragment.
细胞定位	Cell membrane.


图片



Human IGF-I R on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.




Immobilized Human IGF-I R at 5 µg/mL (100 µL/well) can bind Biotinylated Human IGF-II, Fc Tag with a linear range of 0.156-5 µg/mL.



Immobilized Human IGF-I R at 5 µg/mL (100 µL/well) can bind Human IGF-I, Fc Tag with a linear range of 0.04-0.313 µg/mL.

Functional Studies - Recombinant human IGF1 Receptor protein (ab155622)



SDS-PAGE analysis of reduced ab155622 stained overnight with Coomassie Blue.

SDS-PAGE - Recombinant human IGF1 Receptor protein (ab155622)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors