abcam

Product datasheet

Recombinant human RANTES protein ab9680

1 References

描述

产品名称 重组人RANTES蛋白

纯**度** > 98 % SDS-PAGE.

Sterile filtered Greater than 98% pure by HPLC analyses. Endotoxin level is less than 0.1 ng per g

(1EU/g).

表达系统 Escherichia coli

蛋白长度 Full length protein

无动物成分 No

性质 Recombinant

种属Human预测分子量10 kDa

技术指标

Our **Abpromise guarantee** covers the use of **ab9680** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用 Functional Studies

SDS-PAGE

形式 Lyophilized

补充说明 The biological activity of this product is determined by its ability to chemoattract human blood

monocytes using a concentration of 1.0-10.0 ng/ml.

制备和贮存

稳定性和存储 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

This product is an active protein and may elicit a biological response in vivo, handle with caution.

复溶 Please reconstitute in 200ul sterile water.

常规信息

1

功能 Chemoattractant for blood monocytes, memory T-helper cells and eosinophils. Causes the

release of histamine from basophils and activates eosinophils. Binds to CCR1, CCR3, CCR4 and CCR5. One of the major HIV-suppressive factors produced by CD8+ T-cells. Recombinant RANTES protein induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV). The processed form RANTES(3-68) acts as a natural chemotaxis inhibitor and is a more potent inhibitor of HIV-1-infection. The second processed form

RANTES(4-68) exhibits reduced chemotactic and HIV-suppressive activity compared with

RANTES(1-68) and RANTES(3-68) and is generated by an unidentified enzyme associated with

monocytes and neutrophils.

组织特异性 T-cell and macrophage specific.

序列相似性 Belongs to the intercrine beta (chemokine CC) family.

翻译后修饰 N-terminal processed form RANTES(3-68) is produced by proteolytic cleavage, probably by

DPP4, after secretion from peripheral blood leukocytes and cultured sarcoma cells. The identity of the O-linked saccharides at Ser-27 and Ser-28 are not reported in

PubMed:1380064. They are assigned by similarity.

细胞定位 Secreted.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

Guarantee only valid for products bought direct from Abcam or one of our authorized distributors