

Product datasheet

Recombinant Human TCTEX-1 protein ab106872

1 图像

描述

产品名称	重组人TCTEX-1蛋白
纯度	> 95 % SDS-PAGE. ab106872 is purified using conventional chromatography techniques.
表达系统	Escherichia coli
Accession	<u>P63172</u>
蛋白长度	Full length protein
无动物成分	No
性质	Recombinant
种属	Human
序列	MGSSHHHHHSSGLVPRGSHMEDYQAAEETAFFVVDEVSNI VKEAIESAIG GNAYQHSKVNQWTTNVVEQTLSQLTKLGKPFKYIVTCVIMQK NGAGLHTA SSCFWDSSTDGSCVTRWENKTMYSIVSAFGLSI
预测分子量	15 kDa including tags
氨基酸	1 to 113
标签	His tag N-Terminus

技术指标

Our **Abpromise guarantee** covers the use of **ab106872** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	SDS-PAGE Mass Spectrometry
质谱法	MALDI-TOF
形式	Liquid
补充说明	This product was previously labelled as DYNLT1

制备和贮存

稳定性和存储

Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

pH: 8.00

Constituents: 0.0154% DTT, 0.316% Tris HCl, 30% Glycerol (glycerin, glycerine), 0.58% Sodium chloride

常规信息

功能

Acts as one of several non-catalytic accessory components of the cytoplasmic dynein 1 complex that are thought to be involved in linking dynein to cargos and to adapter proteins that regulate dynein function. Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. Binds to transport cargos and is involved in apical cargo transport such as rhodopsin-bearing vesicles in polarized epithelia. Is involved in intracellular targeting of D-type retrovirus gag polyproteins to the cytoplasmic assembly site. May also be a accessory component of axonemal dynein.

Plays a role in neuronal morphogenesis; the function is independent of cytoplasmic dynein and seems to be coupled to regulation of the actin cytoskeleton by enhancing Rac1 activity. The function in neurogenesis may be regulated by association with a G-protein beta-gamma dimer. May function as a receptor-independent activator of heterotrimeric G-protein signaling; the activation appears to be independent of a nucleotide exchange. Plays a role in regulating neurogenesis; inhibits the genesis of neurons from precursor cells during cortical development presumably by antagonizing ARHGEF2. Involved in the regulation of mitotic spindle orientation.

组织特异性

Expressed in heart, placenta, skeletal muscle kidney, pancreas, spleen, prostate, testis, ovary, ileum and colon. Expressed in lung endothelial and smooth muscle cells (at protein level).

序列相似性

Belongs to the dynein light chain Tctex-type family.


翻译后修饰

Phosphorylated by BMPR2; the phosphorylation is abolished by BMPR2 mutations in exon 12 which lead to truncated forms of BMPR2 and which are linked to primary pulmonary hypertension (PPH1) [MIM:178600]. The phosphorylation status is proposed to regulate the association with the cytoplasmic dynein complex and may have role in cytoplasmic dynein cargo release.

细胞定位

Golgi apparatus. Cytoplasm. Cytoplasm > cytoskeleton > spindle. Localizes to mitotic spindles.

图片



15% SDS-PAGE analysis of 3µg ab106872.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors