

# Recombinant mouse TNF alpha protein (Active) ab9740

## 7 References

### 描述

产品名称	重组小鼠TNF alpha蛋白(Active)
生物活性	The <b>ED<sub>50</sub></b> , as determined by the cytolysis of murine L929 cells in the presence of actinomycin D, is $\leq 0.1$ ng/ml, corresponding to a specific activity of $\geq 1 \times 10^7$ units/mg.
纯度	$\geq 98$ % SDS-PAGE. $\geq 98$ % HPLC analyses. Sterile filtered.
内毒素水平	$< 1.000$ Eu/ $\mu$ g
表达系统	Escherichia coli
Accession	<b><u>P06804</u></b>
蛋白长度	Full length protein
无动物成分	No
性质	Recombinant
种属	Mouse
序列	MLRSSSQNSS DKPVAHVVAN HQVEEQLEWL SQRANALLAN GMDLKDNQLV VPADGLYLVY SQVLFKGQGC PDYVLLTHTV SRFAISYQEK VNLLSAVKSP CPKDTPEGAE LKPWYEPIYL GGVFQLEKGD QLSAEVNLPK YLDFAESGQV YFGVIAL
预测分子量	17 kDa
氨基酸	80 to 235

### 技术指标

Our **Abpromise guarantee** covers the use of **ab9740** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

应用	HPLC Functional Studies SDS-PAGE
形式	Lyophilized

### 制备和贮存

<b>稳定性和存储</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.  This product is an active protein and may elicit a biological response in vivo, handle with caution.
<b>复溶</b>	Reconstitute this product in 200ul water.
<b>常规信息</b>	
<b>功能</b>	Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation.
<b>疾病相关</b>	Genetic variations in TNF are a cause of susceptibility psoriatic arthritis (PSORAS) [MIM:607507]. PSORAS is an inflammatory, seronegative arthritis associated with psoriasis. It is a heterogeneous disorder ranging from a mild, non-destructive disease to a severe, progressive, erosive arthropathy. Five types of psoriatic arthritis have been defined: asymmetrical oligoarthritis characterized by primary involvement of the small joints of the fingers or toes; asymmetrical arthritis which involves the joints of the extremities; symmetrical polyarthritis characterized by a rheumatoidlike pattern that can involve hands, wrists, ankles, and feet; arthritis mutilans, which is a rare but deforming and destructive condition; arthritis of the sacroiliac joints and spine (psoriatic spondylitis).
<b>序列相似性</b>	Belongs to the tumor necrosis factor family.
<b>翻译后修饰</b>	The soluble form derives from the membrane form by proteolytic processing. The membrane form, but not the soluble form, is phosphorylated on serine residues. Dephosphorylation of the membrane form occurs by binding to soluble TNFRSF1A/TNFR1. O-glycosylated; glycans contain galactose, N-acetylgalactosamine and N-acetylneuraminic acid.
<b>细胞定位</b>	Secreted and Cell membrane.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors

