abcam

Product datasheet

Biotinylated Goat anti-Mouse & Rabbit IgG (H+L) (Ready To Use) ab64257

10 References 1 图像

概述

产品名称 Biotinylated山羊抗小鼠&兔lgG (H+L) (Ready To Use)

经测试应用 适用于: IHC-P

常规说明 Staining protocol

Deparaffinize and rehydrate formalin-fixed paraffin-embedded tissue section.

Add enough drops of an appropriate hydrogen peroxide block to cover the sections.

Incubate for 10 minutes. Wash 2 times in buffer.

Perform appropriate pretreatment if required. Wash 3 times in buffer.

Apply an appropriate protein block and incubate for 5 minutes at room temperature to block nonspecific background staining. Wash 1 time in buffer.

Apply primary antibody and incubate according to manufacturer's protocol.

Wash 4 times in buffer. Apply Biotinylated Goat anti Mouse & Rabbit lgG(H+L) solution (ab64257) and incubate for 10 minutes at room temperature. Wash 4 times in buffer.

Apply enzyme labeled streptavidin and incubate according to manufacturer's protocol.

Rinse 4 times in buffer. Place slide in appropriate chromogenic substrate and incubate until desired reaction is achieved. Rinse 4 times in buffer.

Add enough drops of Hematoxylin to cover the section. Incubate for 1 minute.

Rinse 7-8 times in tap water. Add Mounting Medium to cover the section.

This reagent is intended to be used in a labeled streptavidin-biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody which reacts with the primary antibody, enzyme-labeled streptavidin, and substrate-chromogen.

性能

形式 Liquid

存放说明 Store at +4°C.

存储溶液 pH: 7.2

Preservative: 0.1% Sodium azide

Constituent: PBS

Reagent说明

This reagent is intended to be used in a labeled streptavidin-biotin immunoenzymatic antigen detection system. This technique involves the sequential incubation of the specimen with an unconjugated primary antibody specific to the target antigen, a biotinylated secondary antibody

1

which reacts with the primary antibody, enzyme-labeled streptavidin, and substrate-chromogen.

细胞定位 Secreted 细胞定位 Secreted


应用

The Abpromise guarantee Abpromise™承诺保证使用ab64257于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应 用 | Ab评论 | 说 明 |
|------------|------|-------------------------------------|
| IHC-P | | Use at an assay dependent dilution. |

图片



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Biotinylated Goat anti-Mouse & Rabbit IgG (H+L) (Ready To Use) (ab64257)

Human colon carcinoma stained with anti-p53 antibody using Biotinylated Goat anti Mouse & Rabbit lgG (H+L) (Ready To Use) (ab64257).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

| • | Guarantee only valid for products bought direct from Abcam or one of our authorized distributors | | | | |
|---|--|---|--|--|--|
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | | | | |
| | | 3 | | | |