# abcam

## Product datasheet

## Glutathione Sepharose® ab193267

## 2 References

1	k	100	,	•	b
7	N	н	٠.	ú	h

产品名称

Glutathione Sepharose®

产品概述

Contents:

Supplied as a 50 % slurry in 20 % Ethanol / PBS; > 3 mg (10  $\mu$ mol) glutathione per mL Sepharose<sup>®</sup> beads.

#### Features:

The Glutathione-S-Transferase (GST) gene fusion system has been widely used for the overexpression of foreign genes in E. Coli and other expression systems. The GST-tagged protein can be easily purified from cell lysate or media by affinity chromatography using glutathione Sepharose<sup>®</sup> beads. Abcam's Glutathione Sepharose<sup>®</sup> is designed for the purification of GST fusion proteins and other glutathione binding proteins. This formulation exhibits excellent binding capacity, high flow rate, no significant loss of the glutathione ligand and a pH stability range of 2-10.

These beads are for use in column purification. If used in batch purification, we recommend not exceeding  $150 \times g$  when centrifuging.

Store beads at 4°C.

The beads may be damaged above 40°C.

DO NOT FREEZE.

Wash beads 3 times with 3x bead volume of desired buffer before use.

### Applications:

Purification of GST-fusion proteins or other glutathione-binding proteins.

Sepharose is a registered trademark of GE Healthcare

**说明** This product is manufactured by BioVision, an Abcam company and was previously called 6555

Glutathione Sepharose. 6555-10 is the same size as the 10 ml size of ab193267.

Glutathione Sepharose<sup>®</sup> is prepared by covalently coupling glutathione to epoxy-activated 4% cross-linked Sepharose<sup>®</sup> beads to form a stable thioether linkage. The coupling was optimized to give a high binding capacity, which could be greater than 5 mg of Glutathione-S-transferase

(GST) per ml of wet gel.

经测试应用 适用于: Purification

#### 性能

**存放说明** Store at +4°C. Please refer to protocols.

组 <b>件</b>	1 ml	10 ml	50 ml
Glutathione Sepharose®	1 x 1ml	1 x 10ml	1 x 50ml

## 应用

## The Abpromise guarantee Abpromise™承诺保证使用ab193267于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应 <b>用</b>	Ab评论	说明
Purification		Use at an assay dependent concentration.  Purification of GST-fusion proteins or other glutathione-binding proteins.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

## Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors