

Goat Anti-Mouse IgG H&L (FITC) ab6785

★★★★★ [1 Abreviews](#) [510 References](#) [4 图像](#)

概述

产品名称	山羊抗小鼠IgG H&L (FITC)
宿主	Goat
靶标种属	Mouse
经测试应用	适用于: IHC (PFA fixed), Immunomicroscopy, Flow Cyt, IHC-P, IHC-Fr, ICC/IF, ELISA, WB
免疫原	mouse whole molecule
偶联物	FITC. Ex: 493nm, Em: 528nm

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	pH: 6.50 Preservative: 0.01% Sodium azide Constituents: 0.42% Tripotassium orthophosphate, 0.87% Sodium chloride, 1% BSA
纯度	Affinity purified
纯化说明	This product was prepared from monospecific antiserum by immunoaffinity chromatography using mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities.
共轭说明	Diaminotriazinylaminofluorescein (DTAF) (Molecular Weight 530 daltons) Absorption Wavelength: 495 nm Emission Wavelength: 528 nm Fluorochrome/Protein Ratio: 3.3 moles FITC per mole of Goat IgG
克隆	多克隆
同种型	IgG


应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab6785于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC (PFA fixed)		1/1000.
Immunomicroscopy		Use at an assay dependent concentration.
Flow Cyt		1/500 - 1/2500. PubMed: 17989178
IHC-P		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration. PubMed: 15716365For PFA fixed tissue see PMID 15716365.
ICC/IF	★★★★★ (1)	1/1000 - 1/5000. (PMID16006199)
ELISA		1/10000 - 1/50000.
WB		1/10000. PubMed: 17217624


图片



Immunocytochemistry/ Immunofluorescence - Goat
Anti-Mouse IgG H&L (FITC) (ab6785)


This image is courtesy of an Abreview submitted by J Chai.

ab8978 staining vimentin in human colon fibroblasts by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with methanol, permeabilized with 0.1% Triton X-100 and blocked with Dako serum free protein blocker for 20 minutes at 28°C. Samples were incubated with primary antibody (1/100 in Dako antibody diluent) for 2 hours at 28°C. A FITC-conjugated goat anti-mouse IgG H&L (ab6785) (1/800) was used as the secondary antibody.




Flow Cytometry - Goat Anti-Mouse IgG H&L (FITC)
(ab6785)

Ab6785 was used at dilution 1/100 with the primary antibody [**ab52455**](#) in Flow Cyt. See the review on [**ab52455**](#).



Immunocytochemistry/ Immunofluorescence - Goat
Anti-Mouse IgG H&L (FITC) (ab6785)

Ab6785 was used at dilution 1/2000 with the primary antibody [**ab1220**](#) in ICC/IF. See the review on [**ab1220**](#).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Goat Anti-Mouse IgG H&L (FITC) (ab6785)

Ab6785 was used at dilution 1/300 with the primary antibody **ab6002** in IHC-P. See the review on **ab6002**.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors