

Goat Anti-Rabbit IgG H&L (HRP polymer) ab214880

★★★★★ [3 Abreviews](#) [44 References](#) [2 图像](#)

概述

产品名称	山羊抗兔IgG H&L (HRP polymer)
宿主	Goat
靶标种属	Rabbit
特异性	Based on IEP, ab214880 reacts with heavy (gamma) chains on rabbit IgG, light chains on all rabbit immunoglobulins. No cross-reactivity is observed to non-immunoglobulin rabbit serum proteins, serum proteins from human, mouse or rat.
经测试应用	适用于: IHC-P, Immunoelectrophoresis
免疫原	Full length protein corresponding to Rabbit IgG. (Purified Rabbit IgG, whole molecule).
偶联物	HRP polymer

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	Preservative: 0.1% Proclin 300 Constituents: 1% BSA, 98% PBS
纯度	Affinity purified
纯化说明	ab214880 was affinity purified using solid phase Rabbit IgG. Purity >95% by SDS-PAGE.
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab214880于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (3)	Use at an assay dependent concentration.

应用	Ab评论	说明
Immunoelectrophoresis		Use at an assay dependent concentration. This product is provided in a pre-diluted, ready-to-use format. Incubate 150 µl per slide for 1 hour.

图片

B

K14.GFP+

K14.GFP-

Ki67

CC3

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880)

Sonzogni et al PLoS Biol. 2018 Jun 20;16(6):e2004049. doi: 10.1371/journal.pbio.2004049. eCollection 2018 Jun. Fig S2. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

Representative images IHC for Ki67 (upper panel) and CC3 (lower panel); scale bar 50 µm.

The paraffin sections were deparaffinized and rehydrated, followed by antigen retrieval using sodium citrate buffer (pH 6). The sections were then incubated with 3% hydrogen peroxide to block endogenous peroxidase activity for 10 minutes at room temperature. After 3 washes with TBS, the sections were incubated with 2% bovine serum albumin for 1 hour at room temperature.

Slides were then incubated separately with rabbit anti-Ki67 and rabbit anti-cleaved caspase 3 overnight at 4°C. The slides were washed 3 times and incubated with goat anti-rabbit HRP polymer (1:1, Abcam ab214880) for 2 hours at room temperature. Samples were washed thrice and then developed, counterstained with hematoxylin and then dehydrated before mounting the slides with permount.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880)

Immunohistochemical analysis of ER/PR positive breast cancer tissue stained for ER and detected with ab214880.

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors