

Anti-PYK2 antibody [YE353] ab32571

敲除验证
重组
RabMAb

[29 References](#)
[12 图像](#)

概述

产品名称	Anti-PYK2抗体[YE353]
描述	兔单克隆抗体[YE353] to PYK2
宿主	Rabbit
特异性	This antibody recognizes PYK2. It does not cross react with other FAK family members.
经测试应用	适用于: WB, IHC-P, ICC/IF 不适用于: IP
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide within Human PYK2 aa 1-100 (N terminal). The exact sequence is proprietary. Database link: Q14289
阳性对照	WB: Ramos, Jurkat and RAW264.7 cell lysates and mouse and rat brain tissue lysates. IHC-P: Human, mouse and rat cerebral cortex tissues. ICC/IF: HeLa and PC12 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol, 0.05% BSA
纯度	Protein A purified
克隆	单克隆

克隆编号 YE353
同种型 IgG

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用 ab32571 于以下的经测试应用

“应用说明”部分下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/2000. Detects a band of approximately 116 kDa (predicted molecular weight: 116 kDa). For unpurified use at 1/1000 - 1/5000.
IHC-P		1/300. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See IHC antigen retrieval protocols . For unpurified use at 1/250 - 1/500.
ICC/IF		1/60. For unpurified use at 1/100.

应用说明 Is unsuitable for IP.

靶标


功能 Involved in calcium induced regulation of ion channel and activation of the map kinase signaling pathway. May represent an important signaling intermediate between neuropeptide activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. Interacts with the SH2 domain of Grb2. May phosphorylate the voltage-gated potassium channel protein Kv1.2. Its activation is highly correlated with the stimulation of c-Jun N-terminal kinase activity. Involved in osmotic stress-dependent SNCA 'Tyr-125' phosphorylation. In concert with SRC, plays an important role in osteoclastic bone resorption. Both the formation of a SRC-PTK2B complex, and SRC kinase activity are necessary for this function. The Tyr-402 phosphorylated form serves as a docking site for SRC and is important for the organization of the osteoclast actin cytoskeleton and attachment sites and for bone resorption.

组织特异性 Most abundant in the brain, with highest levels in amygdala and hippocampus. Low levels in kidney. Also expressed in spleen and lymphocytes.

序列相似性 Belongs to the protein kinase superfamily. Tyr protein kinase family. FAK subfamily. Contains 1 FERM domain. Contains 1 protein kinase domain.

翻译后修饰 Phosphorylated on tyrosine residues in response to various stimuli that elevate the intracellular calcium concentration, as well as by PKC activation. Recruitment by nephrocystin to cell matrix adhesions initiates Tyr-402 phosphorylation. In monocytes, adherence to substrata is required for tyrosine phosphorylation and kinase activation. Angiotensin II, thapsigargin and L-alpha-lysophosphatidic acid (LPA) also induce autophosphorylation and increase kinase activity.

细胞定位 Cytoplasm. Cell membrane. Interaction with nephrocystin induces the membrane-association of the kinase.



Western blot - Anti-PYK2 antibody [YE353] (ab32571)

Lane 1: Wild-type HAP1 whole cell lysate (20 µg)


Lane 2: PYK2 knockout HAP1 whole cell lysate (20 µg)

Lane 3: Jurkat whole cell lysate (20 µg)

Lane 4: Hu brain whole cell lysate (20 µg)


Lanes 1 - 4: Merged signal (red and green). Green - ab32571 observed at 125 kDa. Red - loading control, **ab8245**, observed at 37 kDa.

ab32571 was shown to specifically recognize PYK2 in wild-type HAP1 cells along with additional cross reactive bands. No band was observed when PYK2 knockout samples were examined. Wild-type and PYK2 knockout samples were subjected to SDS-PAGE. Ab32571 and **ab8245** (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-PYK2 antibody [YE353] (ab32571)

Immunocytochemistry/Immunofluorescence analysis of PC12 cells labelling PYK2 with unpurified ab32571 at a 1/100 dilution.



Western blot - Anti-PYK2 antibody [YE353]
(ab32571)

All lanes : Anti-PYK2 antibody [YE353] (ab32571) at 1/10000 dilution (purified)

Lane 1 : Ramos cell lysate

Lane 2 : Jurkat cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 116 kDa

Observed band size: 116 kDa

Blocking buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm /TBST.



Western blot - Anti-PYK2 antibody [YE353]
(ab32571)

Anti-PYK2 antibody [YE353] (ab32571) at 10000 cells (purified) + RAW264.7 cell lysate at 20 µg

Secondary


Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 116 kDa

Observed band size: 116 kDa

Blocking buffer and concentration: 5% NFDm/TBST.

Diluting buffer and concentration: 5% NFDm /TBST.



Western blot - Anti-PYK2 antibody [YE353]
(ab32571)

All lanes : Anti-PYK2 antibody [YE353] (ab32571) at 1/2000 dilution (purified)

Lane 1 : Mouse brain tissue lysate

Lane 2 : Rat brain tissue lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 116 kDa

Observed band size: 116 kDa

Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM /TBST.




Western blot - Anti-PYK2 antibody [YE353]
(ab32571)

Anti-PYK2 antibody [YE353] (ab32571) at 1/5000 dilution (unpurified) + Jurkat cell lysate


Predicted band size: 116 kDa

Observed band size: 116 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cerebral cortex tissue labelling PYK2 with purified ab32571 at 1/300. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. [ab97051](#), a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PYK2 antibody [YE353] (ab32571)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse cerebral cortex tissue labelling PYK2 with purified ab32571 at 1/300. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. [ab97051](#), a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PYK2 antibody [YE353] (ab32571)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of rat cerebral cortex tissue labelling PYK2 with purified ab32571 at 1/300. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PYK2 antibody [YE353] (ab32571)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of normal human brain tissue labelling PYK2 with unpurified ab32571 at a 1/250 dilution.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PYK2 antibody [YE353] (ab32571)



Immunocytochemistry/ Immunofluorescence - Anti-PYK2 antibody [YE353] (ab32571)

Immunocytochemistry/Immunofluorescence analysis of HeLa cells labelling PYK2 with purified ab32571 at 1/60. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain.

Control: primary antibody (1/100) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Why choose a recombinant antibody?

 Research with confidence Consistent and reproducible results	 Long-term and scalable supply Recombinant technology
 Success from the first experiment Confirmed specificity	 Ethical standards compliant Animal-free production

Anti-PYK2 antibody [YE353] (ab32571)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors